

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/29, 15/54, 15/55, 15/61, 15/82, C07K 14/415, A01H 5/00	A2	(11) International Publication Number: WO 00/12716 (43) International Publication Date: 9 March 2000 (09.03.00)
(21) International Application Number: PCT/US99/20051 (22) International Filing Date: 31 August 1999 (31.08.99) (30) Priority Data: 60/098,986 2 September 1998 (02.09.98) : US (71) Applicant: PIONEER HI-BRED INTERNATIONAL, INC. [US/US]; 800 Capital Square, 400 Locust Street, Des Moines, IA 50309 (US). (72) Inventors: MAHAJAN, Pramod, B.; 8029 Brookview Drive, Urbandale, IA 50322 (US). SHI, Jinrui; 6022 Greendale Circle, Apartment 205, Johnston, IA 50131 (US). (74) Agents: YATES, Michael, E. et al.; Darwin Building, 7100 N.W. 62nd Avenue, Johnston, IA 50131-1000 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: MAIZE Ku70 ORTHOLOGUE AND USES THEREOF (57) Abstract The invention provides isolated maize Ku70 nucleic acids and their encoded proteins. The present invention provides methods and compositions relating to altering maize Ku70 concentration and/or composition of plants. The invention further provides recombinant expression cassettes, host cells, transgenic plants, and antibody compositions.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Maize Ku70 Orthologue and Uses Thereof

TECHNICAL FIELD

The present invention relates generally to plant molecular biology. More specifically, it relates to nucleic acids and methods for modulating their expression in plants.

BACKGROUND OF THE INVENTION

Cellular DNA undergoes double strand breakage during the course of many physiological events as well as in response to a variety of environmental insults (1, 2). Left unrepaired, such double strand breaks (DSBs) lead to mutations that may prove lethal to the organism. Therefore, these DSBs are repaired promptly via two independent pathways: i) homologous recombination ii) non-homologous end joining (1,2). The first pathway involves a series of very specific biochemical reactions catalyzed by a complex of cellular proteins (3). Due to the large number of proteins involved in this complex, it is referred to as a 'recombinosome' (4). This pathway is the dominant mode of DSB repair in lower eukaryotes such as yeast (2).

The non-homologous end-joining pathway is the major route of DSB repair in higher eukaryotes (1,2). This pathway is also catalyzed by a group of cellular proteins. This group contains, in addition to hitherto unidentified factors, some well-characterized enzymes such as DNA ligases, Poly (ADP-Ribose) Polymerase [PADPRP], and DNA - dependent Protein Kinase [DNA-PK] (5,6). These enzymes have been studied in detail using lower as well as higher vertebrate systems including mammals. Both PADPRP and DNA-PK have been shown to be activated by DNA ends. Moreover, these two enzymes also bind DNA ends (5,6). While PADPRP is a single polypeptide of ~115 kDa (7), DNA-PK exists as a complex of two subunits (8,9). The catalytic subunit [DNA-PK_{cat}] is composed of a single polypeptide of ~450kDa. It is a serine-threonine type of protein kinase that phosphorylates a variety of nuclear enzymes, transcription factors and oncogenes (9). However, DNA-PK_{cat} by itself does not bind DNA. The non-catalytic subunit of DNA-PK is a heterodimer composed of 70kDa and 86kDa proteins. The non-catalytic subunit acts as a regulator of DNA-PK_{cat} by virtue of its' ability to bind to DNA ends, thereby recruiting the catalytic subunit to the site of DSBs (8,9).

- 2 -

Although enzymology of DNA-PK_{cs} has been investigated extensively (8), its biological function was identified only recently (10). Availability of the full length cDNA sequence of mammalian DNA-PK_{cs} allowed identification of this protein as a member of the phosphatidylinositol 3-kinase (PI kinase) gene family. While most members of this family are lipid kinases, a small number of proteins forming a subfamily specifically phosphorylate proteins. Members of this subfamily are known as PI-K related kinases and include the ATM protein, Tel1p, Tor1p, Tor2p, FRAP, Rad3p, Mec1p and Mei41 (10). In addition to their structural and biochemical similarities, members of this subfamily also appear to share a common biological function. They are all involved in repair of DNA that is damaged in response to a variety of genetic, physiological or environmental events (10). Although several members of this subfamily have been cloned from animals, no known information on plant DNA-PK_{cs} is available in the literature.

The non-catalytic subunit of DNA-PK appear to be identical to previously well characterized mammalian Ku proteins (8). The Ku complex, also a heterodimer of 70kDa and 86kDa proteins, was shown to be a nuclear DNA-binding autoantigen (12,13). Patients diagnosed of a variety of autoimmune diseases have been known to develop antibodies to Ku proteins (14). Further biochemical analysis has established that Ku binds with strong affinity to DNA ends, stem-loop structures, DNA bubbles, or transitions between double stranded DNA and two single strands (15). Subsequent to binding to the ends, Ku molecules can translocate along the DNA, such that three or more molecules can bind to the linear DNA fragment. Both components of Ku have a DNA dependent ATPase activity and an ATP dependent helicase activity (15). Recently, Yoo and Dynan have also demonstrated RNA binding activity of the Ku protein (16).

Recent genetic studies using rodent cell lines defective in DNA strand break repair have provided the important link between Ku protein, DNA-PK and DSB repairs during DNA replication, repair and recombination (16). Boulton & Jackson have shown that the yeast Ku70 potentiates illegitimate DNA DSB repair and serves as a barrier to error-prone DNA repair pathways (17). Studies with mutant rodent cell lines have clearly shown that Ku proteins are required for the V (D) J DNA recombination (18) and immunoglobulin isotype switching (19). Components of DNA-PK are also involved in the non-homologous end-joining pathway in telomeric length maintenance and telomere silencing (20) as well as telomere integrity (21). Ramsden & Gellert have recently

- 3 -

observed that Ku protein stimulates DNA end joining by mammalian DNA ligases and proposed a direct role for Ku in DSB repair (22). A role for Ku protein in modulation of heat shock response (23), and hyperthermic radiosensitization (24) has also been advocated. As discussed above, recent studies have established the role of DNA-PK components in various cellular processes involving DSB. During the course of these investigations, Ku orthologues have been cloned from human (25, 26), mouse (27), *Drosophila melanogaster* (28), *Rhipicephalus appendiculatus* (29) and *Caenorhabditis elegans* (30). Interestingly, Ku orthologues have also been reported in *Saccharomyces cerevisiae* (31-33).

Control of homologous recombination or non-homologous end joining by modulating Ku provides the means to modulate the efficiency which heterologous nucleic acids are incorporated into the genomes of a target plant cell. Control of these processes has important implications in the creation of novel recombinantly engineered crops such as maize. The present invention provides this and other advantages.

15

SUMMARY OF THE INVENTION

Generally, it is the object of the present invention to provide nucleic acids and proteins relating to the maize Ku70 orthologue. It is an object of the present invention to provide: 1) antigenic fragments of the proteins of the present invention; 2) transgenic plants comprising the nucleic acids of the present invention; 3) methods for modulating, in a transgenic plant, the expression of the nucleic acids of the present invention.

Therefore, in one aspect, the present invention relates to an isolated nucleic acid comprising a member selected from the group consisting of (a) a polynucleotide having a specified sequence identity to a polynucleotide encoding a polypeptide of the present invention, wherein the polypeptide when presented as an immunogen elicits the production of an antibody which is specifically reactive to the polypeptide; (b) a polynucleotide which is complementary to the polynucleotide of (a); and (c) a polynucleotide comprising a specified number of contiguous nucleotides from a polynucleotide of (a) or (b). The isolated nucleic acid can be DNA.

In another aspect, the present invention relates to recombinant expression cassettes, comprising a nucleic acid as described, *supra*, operably linked to a promoter. In some embodiments, the nucleic acid is operably linked in antisense orientation to the promoter.

- 4 -

In another aspect, the present invention is directed to a host cell transfected with the recombinant expression cassette as described, *supra*. In some embodiments, the host cell is a sorghum (*Sorghum bicolor*) or maize (*Zea mays*) cell.

In a further aspect, the present invention relates to an isolated protein comprising
5 a polypeptide having a specified number of contiguous amino acids encoded by the isolated nucleic acid referred to, *supra*.

In another aspect, the present invention relates to an isolated nucleic acid comprising a polynucleotide of specified length which selectively hybridizes under stringent conditions to a nucleic acid of the present invention, or a complement thereof.

10 In some embodiments, the isolated nucleic acid is operably linked to a promoter.

In yet another aspect, the present invention relates to an isolated nucleic acid comprising a polynucleotide, the polynucleotide having a specified sequence identity to an identical length of a nucleic acid of the present invention or a complement thereof.

In another aspect, the present invention relates to an isolated nucleic acid
15 comprising a polynucleotide having a sequence of a nucleic acid amplified from a *Zea mays* nucleic acid library using at least two primers or their complements, one of which selectively hybridizes under stringent conditions to a locus of the nucleic acid comprising the 5' terminal coding region and the other primer selectively hybridizing, under stringent conditions, to a locus of the nucleic acid comprising the 3' terminal coding
20 region, and wherein both primers selectively hybridize within the coding region. In some embodiments, the nucleic acid library is a cDNA library.

In another aspect, the present invention relates to a recombinant expression cassette comprising a nucleic acid amplified from a library as referred to *supra*, wherein the nucleic acid is operably linked to a promoter. In some embodiments, the present
25 invention relates to a host cell transfected with this recombinant expression cassette. In some embodiments, the present invention relates to a protein of the present invention which is produced from this host cell.

In an additional aspect, the present invention is directed to an isolated nucleic acid comprising a polynucleotide encoding a polypeptide wherein: (a) the polypeptide
30 comprises a specified number of contiguous amino acid residues from a first polypeptide of the present invention, wherein the polypeptide, when presented as an immunogen, elicits the production of an antibody which specifically binds to said first polypeptide; (b)

- 5 -

the polypeptide does not bind to antisera raised against the first polypeptide which has been fully immunosorbed with the first polypeptide; (c) the polypeptide has a molecular weight in non-glycosylated form within a specified percentage of the first polypeptide.

5 In a further aspect, the present invention relates to a heterologous promoter operably linked to a non-isolated polynucleotide of the present invention, wherein the polypeptide is encoded by a nucleic acid amplified from a nucleic acid library.

 In yet another aspect, the present invention relates to a transgenic plant comprising a recombinant expression cassette comprising a plant promoter operably
10 linked to any of the isolated nucleic acids of the present invention. In some embodiments, the transgenic plant is *Zea mays*. The present invention also provides transgenic seed from the transgenic plant.

 In a further aspect, the present invention relates to a method of modulating expression of the genes encoding the proteins of the present invention in a plant,
15 comprising the steps of (a) transforming a plant cell with a recombinant expression cassette comprising a polynucleotide of the present invention operably linked to a promoter; (b) growing the plant cell under plant growing conditions; and (c) inducing expression of the polynucleotide for a time sufficient to modulate expression of the genes in the plant. In some embodiments, the plant is maize. Expression of the genes
20 encoding the proteins of the present invention can be increased or decreased relative to a non-transformed control plant.

Definitions

 Units, prefixes, and symbols may be denoted in their SI accepted form. Unless
25 otherwise indicated, nucleic acids are written left to right in 5' to 3' orientation; amino acid sequences are written left to right in amino to carboxy orientation, respectively. Numeric ranges are inclusive of the numbers defining the range and include each integer within the defined range. Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the
30 IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes. The terms defined below are more fully defined by reference to the specification as a whole.

By "amplified" is meant the construction of multiple copies of a nucleic acid sequence or multiple copies complementary to the nucleic acid sequence using at least one of the nucleic acid sequences as a template. Amplification systems include the polymerase chain reaction (PCR) system, ligase chain reaction (LCR) system, nucleic acid sequence based amplification (NASBA, Cingone, Mississauga, Ontario), Q-Beta Replicase systems, transcription-based amplification system (TAS), and strand displacement amplification (SDA). See, e.g., *Diagnostic Molecular Microbiology: Principles and Applications*, D. H. Persing *et al.*, Ed., American Society for Microbiology, Washington, D.C. (1993). The product of amplification is termed an amplicon.

The term "antibody" includes reference to antigen binding forms of antibodies (e.g., Fab, F(ab)₂). The term "antibody" frequently refers to a polypeptide substantially encoded by an immunoglobulin gene or immunoglobulin genes, or fragments thereof which specifically bind and recognize an analyte (antigen). However, while various antibody fragments can be defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized *de novo* either chemically or by utilizing recombinant DNA methodology. Thus, the term antibody, as used herein, also includes antibody fragments such as single chain Fv, chimeric antibodies (i.e., comprising constant and variable regions from different species), humanized antibodies (i.e., comprising a complementarity determining region (CDR) from a non-human source) and heteroconjugate antibodies (e.g., bispecific antibodies).

The term "antigen" includes reference to a substance to which an antibody can be generated and/or to which the antibody is specifically immunoreactive. The specific immunoreactive sites within the antigen are known as epitopes or antigenic determinants. These epitopes can be a linear array of monomers in a polymeric composition - such as amino acids in a protein - or consist of or comprise a more complex secondary or tertiary structure. Those of skill will recognize that all immunogens (i.e., substances capable of eliciting an immune response) are antigens; however some antigens, such as haptens, are not immunogens but may be made immunogenic by coupling to a carrier molecule. An antibody immunologically reactive with a particular antigen can be generated *in vivo* or by recombinant methods such as selection of libraries of recombinant antibodies in phage or similar vectors. See, e.g., Huse *et al.*, *Science* 246: 1275-1281 (1989); and Ward, *et*

al., *Nature* 341: 544-546 (1989); and Vaughan *et al.*, *Nature Biotech.* 14: 309-314 (1996).

As used herein, "antisense orientation" includes reference to a duplex polynucleotide sequence which is operably linked to a promoter in an orientation where the antisense strand is transcribed. The antisense strand is sufficiently complementary to an endogenous transcription product such that translation of the endogenous transcription product is often inhibited.

As used herein, "chromosomal region" includes reference to a length of a chromosome which may be measured by reference to the linear segment of DNA which it comprises. The chromosomal region can be defined by reference to two unique DNA sequences, i.e., markers.

The term "conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or conservatively modified variants of the amino acid sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations" and represent one species of conservatively modified variation. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the nucleic acid. One of ordinary skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine; and UGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide of the present invention is implicit in each described polypeptide sequence and incorporated herein by reference.

As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a "conservatively modified variant" where the alteration results in the substitution of an amino acid with a chemically similar amino

- 8 -

acid. Thus, any number of amino acid residues selected from the group of integers consisting of from 1 to 15 can be so altered. Thus, for example, 1, 2, 3, 4, 5, 7, or 10 alterations can be made. Conservatively modified variants typically provide similar biological activity as the unmodified polypeptide sequence from which they are derived.

- 5 For example, substrate specificity, enzyme activity, or ligand/receptor binding is generally at least 30%, 40%, 50%, 60%, 70%, 80%, or 90% of the native protein for its native substrate. Conservative substitution tables providing functionally similar amino acids are well known in the art.

The following six groups each contain amino acids that are conservative
10 substitutions for one another:

- 1) Alanine (A), Serine (S), Threonine (T);
- 2) Aspartic acid (D), Glutamic acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 15 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

See also, Creighton (1984) *Proteins* W.H. Freeman and Company.

By "encoding" or "encoded", with respect to a specified nucleic acid, is meant comprising the information for translation into the specified protein. A nucleic acid
20 encoding a protein may comprise non-translated sequences (e.g., introns) within translated regions of the nucleic acid, or may lack such intervening non-translated sequences (e.g., as in cDNA). The information by which a protein is encoded is specified by the use of codons. Typically, the amino acid sequence is encoded by the nucleic acid using the "universal" genetic code. However, variants of the universal
25 code, such as are present in some plant, animal, and fungal mitochondria, the bacterium *Mycoplasma capricolum* (*Proc. Natl. Acad. Sci. (USA)*, 82: 2306-2309 (1985)), or the ciliate *Macronucleus*, may be used when the nucleic acid is expressed using these organisms.

When the nucleic acid is prepared or altered synthetically, advantage can be taken
30 of known codon preferences of the intended host where the nucleic acid is to be expressed. For example, although nucleic acid sequences of the present invention may be expressed in both monocotyledonous and dicotyledonous plant species, sequences can be modified to account for the specific codon preferences and GC content preferences of

monocotyledons or dicotyledons as these preferences have been shown to differ (Murray *et al.* Nucl. Acids Res. 17: 477-498 (1989)). Thus, the maize preferred codon for a particular amino acid may be derived from known gene sequences from maize. Maize codon usage for 28 genes from maize plants are listed in Table 4 of Murray *et al.*, *supra*.

- 5 As used herein "full-length sequence" in reference to a specified polynucleotide or its encoded protein means having the entire amino acid sequence of, a native (non-synthetic), endogenous, catalytically active form of the specified protein. A full-length sequence can be determined by size comparison relative to a control which is a native (non-synthetic) endogenous cellular form of the specified nucleic acid or protein.
- 10 Methods to determine whether a sequence is full-length are well known in the art including such exemplary techniques as northern or western blots, primer extension, S1 protection, and ribonuclease protection. See, e.g., *Plant Molecular Biology: A Laboratory Manual*, Clark, Ed., Springer-Verlag, Berlin (1997). Comparison to known full-length homologous (orthologous and/or paralogous) sequences can also be used to
- 15 identify full-length sequences of the present invention. Additionally, consensus sequences typically present at the 5' and 3' untranslated regions of mRNA aid in the identification of a polynucleotide as full-length. For example, the consensus sequence ANNNNAAUGG, where the underlined codon represents the N-terminal methionine, aids in determining whether the polynucleotide has a complete 5' end. Consensus sequences
- 20 at the 3' end, such as polyadenylation sequences, aid in determining whether the polynucleotide has a complete 3' end.

- As used herein, "heterologous" in reference to a nucleic acid is a nucleic acid that originates from a foreign species, or, if from the same species, is substantially modified from its native form in composition and/or genomic locus by deliberate human
- 25 intervention. For example, a promoter operably linked to a heterologous structural gene is from a species different from that from which the structural gene was derived, or, if from the same species, one or both are substantially modified from their original form. A heterologous protein may originate from a foreign species or, if from the same species, is substantially modified from its original form by deliberate human
- 30 intervention.

By "host cell" is meant a cell which contains a vector and supports the replication and/or expression of the expression vector. Host cells may be prokaryotic cells such as *E. coli*, or eukaryotic cells such as yeast, insect, amphibian, or mammalian cells.

Preferably, host cells are monocotyledonous or dicotyledonous plant cells. A particularly preferred monocotyledonous host cell is a maize host cell.

The term "hybridization complex" includes reference to a duplex nucleic acid structure formed by two single-stranded nucleic acid sequences selectively hybridized with each other.

By "immunologically reactive conditions" or "immunoreactive conditions" is meant conditions which allow an antibody, generated to a particular epitope, to bind to that epitope to a detectably greater degree (e.g., at least 2-fold over background) than the antibody binds to substantially all other epitopes in a reaction mixture comprising the particular epitope. Immunologically reactive conditions are dependent upon the format of the antibody binding reaction and typically are those utilized in immunoassay protocols. See Harlow and Lane, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, New York (1988), for a description of immunoassay formats and conditions.

The term "introduced" in the context of inserting a nucleic acid into a cell, means "transfection" or "transformation" or "transduction" and includes reference to the incorporation of a nucleic acid into a eukaryotic or prokaryotic cell where the nucleic acid may be incorporated into the genome of the cell (e.g., chromosome, plasmid, plastid or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (e.g., transfected mRNA).

The terms "isolated" refers to material, such as a nucleic acid or a protein, which is: (1) substantially or essentially free from components which normally accompany or interact with it as found in its naturally occurring environment. The isolated material optionally comprises material not found with the material in its natural environment; or (2) if the material is in its natural environment, the material has been synthetically (non-naturally) altered by deliberate human intervention to a composition and/or placed at a locus in the cell (e.g., genome or subcellular organelle) not native to a material found in that environment. The alteration to yield the synthetic material can be performed on the material within or removed from its natural state. For example, a naturally occurring nucleic acid becomes an isolated nucleic acid if it is altered, or if it is transcribed from DNA which has been altered, by non-natural, synthetic (i.e., "man-made") methods performed within the cell from which it originates. See, e.g., Compounds and Methods for Site Directed Mutagenesis in Eukaryotic Cells, Kmiec, U.S. Patent No. 5,565,350;

In Vivo Homologous Sequence Targeting in Eukaryotic Cells; Zarling *et al.*, PCT/US93/03868. Likewise, a naturally occurring nucleic acid (e.g., a promoter) becomes isolated if it is introduced by non-naturally occurring means to a locus of the genome not native to that nucleic acid. Nucleic acids which are "isolated" as defined
5 herein, are also referred to as "heterologous" nucleic acids.

Unless otherwise stated, the term "maize Ku70 nucleic acid" is a nucleic acid of the present invention and means a nucleic acid comprising a polynucleotide of the present invention (a "maize Ku70 polynucleotide") encoding a maize Ku70 polypeptide. A "maize Ku70 gene" is a gene of the present invention and refers to a non-heterologous
10 genomic form of a full-length maize Ku70 polynucleotide.

As used herein, "localized within the chromosomal region defined by and including" with respect to particular markers includes reference to a contiguous length of a chromosome delimited by and including the stated markers.

As used herein, "marker" includes reference to a locus on a chromosome that
15 serves to identify a unique position on the chromosome. A "polymorphic marker" includes reference to a marker which appears in multiple forms (alleles) such that different forms of the marker, when they are present in a homologous pair, allow transmission of each of the chromosomes in that pair to be followed. A genotype may be defined by use of one or a plurality of markers.

20 As used herein, "nucleic acid" includes reference to a deoxyribonucleotide or ribonucleotide polymer in either single- or double-stranded form, and unless otherwise limited, encompasses known analogues having the essential nature of natural nucleotides in that they hybridize to single-stranded nucleic acids in a manner similar to naturally occurring nucleotides (e.g., peptide nucleic acids).

25 By "nucleic acid library" is meant a collection of isolated DNA or RNA molecules which comprise and substantially represent the entire transcribed fraction of a genome of a specified organism. Construction of exemplary nucleic acid libraries, such as genomic and cDNA libraries, is taught in standard molecular biology references such as Berger and Kimmel, *Guide to Molecular Cloning Techniques, Methods in*
30 *Enzymology*, Vol. 152, Academic Press, Inc., San Diego, CA (Berger); Sambrook *et al.*, *Molecular Cloning - A Laboratory Manual*, 2nd ed., Vol. 1-3 (1989); and *Current Protocols in Molecular Biology*, F.M. Ausubel *et al.*, Eds., Current Protocols, a joint

venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc. (1994 Supplement).

As used herein "operably linked" includes reference to a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. Generally, operably linked means that the nucleic acid sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame.

As used herein, the term "plant" includes reference to whole plants, plant organs (e.g., leaves, stems, roots, etc.), seeds and plant cells and progeny of same. Plant cell, as used herein includes, without limitation, seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores. The class of plants which can be used in the methods of the invention is generally as broad as the class of higher plants amenable to transformation techniques, including both monocotyledonous and dicotyledonous plants. A particularly preferred plant is *Zea mays*.

As used herein, "polynucleotide" includes reference to a deoxyribopolynucleotide, ribopolynucleotide, or analogs thereof that have the essential nature of a natural ribonucleotide in that they hybridize, under stringent hybridization conditions, to substantially the same nucleotide sequence as naturally occurring nucleotides and/or allow translation into the same amino acid(s) as the naturally occurring nucleotide(s). A polynucleotide can be full-length or a subsequence of a native or heterologous structural or regulatory gene. Unless otherwise indicated, the term includes reference to the specified sequence as well as the complementary sequence thereof. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotides" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term polynucleotide as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including *inter alia*, simple and complex cells.

The terms "polypeptide", "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical analogue of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers. The essential nature of such analogues of naturally occurring amino acids is that, when incorporated into a protein, that protein is specifically reactive to antibodies elicited to the same protein but consisting entirely of naturally occurring amino acids. The terms "polypeptide", "peptide" and "protein" are also inclusive of modifications including, but not limited to, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation. Exemplary modifications are described in most basic texts, such as, *Proteins - Structure and Molecular Properties*, 2nd ed., T. E. Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as, for example, those provided by Wold, F., *Post-translational Protein Modifications: Perspectives and Prospects*, pp. 1-12 in *Posttranslational Covalent Modification of Proteins*, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter *et al.*, *Meth. Enzymol.* 182: 626-646 (1990) and Rattan *et al.*, *Protein Synthesis: Posttranslational Modifications and Aging*, *Ann. N.Y. Acad. Sci.* 663: 48-62 (1992). It will be appreciated, as is well known and as noted above, that polypeptides are not always entirely linear. For instance, polypeptides may be branched as a result of ubiquitination, and they may be circular, with or without branching, generally as a result of posttranslation events, including natural processing event and events brought about by human manipulation which do not occur naturally. Circular, branched and branched circular polypeptides may be synthesized by non-translation natural process and by entirely synthetic methods, as well. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. In fact, blockage of the amino or carboxyl group in a polypeptide, or both, by a covalent modification, is common in naturally occurring and synthetic polypeptides and such modifications may be present in polypeptides of the present invention, as well. For instance, the amino terminal residue of polypeptides made in *E. coli* or other cells, prior to proteolytic processing, almost invariably will be N-formylmethionine. During post-translational modification of the peptide, a methionine residue at the NH₂-terminus may be deleted. Accordingly, this invention contemplates the use of both the methionine-containing and the methionine-less

amino terminal variants of the protein of the invention. In general, as used herein, the term polypeptide encompasses all such modifications, particularly those that are present in polypeptides synthesized by expressing a polynucleotide in a host cell.

As used herein "promoter" includes reference to a region of DNA upstream from the start of transcription and involved in recognition and binding of RNA polymerase and other proteins to initiate transcription. A "plant promoter" is a promoter capable of initiating transcription in plant cells. Exemplary plant promoters include, but are not limited to, those that are obtained from plants, plant viruses, and bacteria which comprise genes expressed in plant cells such *Agrobacterium* or *Rhizobium*. Examples of promoters under developmental control include promoters that preferentially initiate transcription in certain tissues, such as leaves, roots, or seeds. Such promoters are referred to as "tissue preferred". Promoters which initiate transcription only in certain tissue are referred to as "tissue specific". A "cell type" specific promoter primarily drives expression in certain cell types in one or more organs, for example, vascular cells in roots or leaves. An "inducible" promoter is a promoter which is under environmental control. Examples of environmental conditions that may effect transcription by inducible promoters include anaerobic conditions or the presence of light. Tissue specific, tissue preferred, cell type specific, and inducible promoters constitute the class of "non-constitutive" promoters. A "constitutive" promoter is a promoter which is active under most environmental conditions.

The term "maize Ku70 polypeptide" is a polypeptide of the present invention and refers to one or more amino acid sequences, in glycosylated or non-glycosylated form. The term is also inclusive of fragments, variants, homologs, alleles or precursors (e.g., preproteins or proproteins) thereof. A "maize Ku70 protein" is a protein of the present invention and comprises a maize Ku70 polypeptide.

As used herein "recombinant" includes reference to a cell or vector, that has been modified by the introduction of a heterologous nucleic acid or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found in identical form within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under-expressed or not expressed at all as a result of deliberate human intervention. The term "recombinant" as used herein does not encompass the alteration of the cell or vector by naturally occurring

- 15 -

events (e.g., spontaneous mutation, natural transformation/transduction/transposition) such as those occurring without deliberate human intervention.

As used herein, a "recombinant expression cassette" is a nucleic acid construct, generated recombinantly or synthetically, with a series of specified nucleic acid elements which permit transcription of a particular nucleic acid in a host cell. The recombinant expression cassette can be incorporated into a plasmid, chromosome, mitochondrial DNA, plastid DNA, virus, or nucleic acid fragment. Typically, the recombinant expression cassette portion of an expression vector includes, among other sequences, a nucleic acid to be transcribed, and a promoter.

The term "residue" or "amino acid residue" or "amino acid" are used interchangeably herein to refer to an amino acid that is incorporated into a protein, polypeptide, or peptide (collectively "protein"). The amino acid may be a naturally occurring amino acid and, unless otherwise limited, may encompass known analogs of natural amino acids that can function in a similar manner as naturally occurring amino acids.

The term "selectively hybridizes" includes reference to hybridization, under stringent hybridization conditions, of a nucleic acid sequence to a specified nucleic acid target sequence to a detectably greater degree (e.g., at least 2-fold over background) than its hybridization to non-target nucleic acid sequences and to the substantial exclusion of non-target nucleic acids. Selectively hybridizing sequences typically have about at least 80% sequence identity, preferably 90% sequence identity, and most preferably 100% sequence identity (i.e., complementary) with each other.

The term "specifically reactive", includes reference to a binding reaction between an antibody and a protein having an epitope recognized by the antigen binding site of the antibody. This binding reaction is determinative of the presence of a protein having the recognized epitope amongst the presence of a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to an analyte having the recognized epitope to a substantially greater degree (e.g., at least 2-fold over background) than to substantially all other analytes lacking the epitope which are present in the sample.

Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, antibodies raised to the polypeptides of the present invention can be selected from to obtain antibodies

specifically reactive with polypeptides of the present invention. The proteins used as immunogens can be in native conformation or denatured so as to provide a linear epitope.

5 A variety of immunoassay formats may be used to select antibodies specifically reactive with a particular protein (or other analyte). For example, solid-phase ELISA immunoassays are routinely used to select monoclonal antibodies specifically immunoreactive with a protein. See Harlow and Lane, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, New York (1988), for a description of immunoassay formats and conditions that can be used to determine selective reactivity.

10 The terms "stringent conditions" or "stringent hybridization conditions" includes reference to conditions under which a probe will hybridize to its target sequence, to a detectably greater degree than other sequences (e.g., at least 2-fold over background). Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing
15 conditions, target sequences can be identified which are 100% complementary to the probe (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleotides in length, preferably less than 500 nucleotides in length.

20 Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of
25 destabilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37°C, and a wash in 1X to 2X SSC (20X SSC = 3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55°C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash
30 in 0.5X to 1X SSC at 55 to 60°C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 0.1X SSC at 60 to 65°C.

Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth and Wahl, *Anal. Biochem.*, 138:267-284 (1984): $T_m = 81.5^{\circ}\text{C} + 16.6 (\log M) + 0.41 (\%GC) - 0.61 (\% \text{ form}) - 500/L$; where M is the molarity of monovalent cations, $\%GC$ is the percentage of guanosine and cytosine nucleotides in the DNA, $\% \text{ form}$ is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The T_m is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. T_m is reduced by about 1°C for each 1% of mismatching; thus, T_m , hybridization and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with $\geq 90\%$ identity are sought, the T_m can be decreased 10°C . Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or 4°C lower than the thermal melting point (T_m); moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9, or 10°C lower than the thermal melting point (T_m); low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or 20°C lower than the thermal melting point (T_m). Using the equation, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45°C (aqueous solution) or 32°C (formamide solution) it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Laboratory Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 "Overview of principles of hybridization and the strategy of nucleic acid probe assays", Elsevier, New York (1993); and *Current Protocols in Molecular Biology*, Chapter 2, Ausubel, *et al.*, Eds., Greene Publishing and Wiley-Interscience, New York (1995).

As used herein, "transgenic plant" includes reference to a plant which comprises within its genome a heterologous polynucleotide. Generally, the heterologous polynucleotide is stably integrated within the genome such that the polynucleotide is

- 18 -

passed on to successive generations. The heterologous polynucleotide may be integrated into the genome alone or as part of a recombinant expression cassette. "Transgenic" is used herein to include any cell, cell line, callus, tissue, plant part or plant, the genotype of which has been altered by the presence of heterologous nucleic acid including those transgenics initially so altered as well as those created by sexual crosses or asexual propagation from the initial transgenic. The term "transgenic" as used herein does not encompass the alteration of the genome (chromosomal or extra-chromosomal) by conventional plant breeding methods or by naturally occurring events such as random cross-fertilization, non-recombinant viral infection, non-recombinant bacterial transformation, non-recombinant transposition, or spontaneous mutation.

As used herein, "vector" includes reference to a nucleic acid used in transfection of a host cell and into which can be inserted a polynucleotide. Vectors are often replicons. Expression vectors permit transcription of a nucleic acid inserted therein.

The following terms are used to describe the sequence relationships between two or more nucleic acids or polynucleotides: (a) "reference sequence", (b) "comparison window", (c) "sequence identity", (d) "percentage of sequence identity", and (e) "substantial identity".

(a) As used herein, "reference sequence" is a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset or the entirety of a specified sequence; for example, as a segment of a full-length cDNA or gene sequence, or the complete cDNA or gene sequence.

(b) As used herein, "comparison window" means includes reference to a contiguous and specified segment of a polynucleotide sequence, wherein the polynucleotide sequence may be compared to a reference sequence and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Generally, the comparison window is at least 20 contiguous nucleotides in length, and optionally can be 30, 40, 50, 100, or longer. Those of skill in the art understand that to avoid a high similarity to a reference sequence due to inclusion of gaps in the polynucleotide sequence a gap penalty is typically introduced and is subtracted from the number of matches.

Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman, *Adv. Appl. Math.* 2: 482 (1981); by the homology alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48: 443 (1970); by the search for similarity method of Pearson and Lipman, *Proc. Natl. Acad. Sci.* 85: 2444 (1988); by computerized implementations of these algorithms, including, but not limited to: CLUSTAL in the PC/Gene program by Intelligenetics, Mountain View, California, GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, Wisconsin, USA; the CLUSTAL program is well described by Higgins and Sharp, *Gene* 73: 237-244 (1988); Higgins and Sharp, *CABIOS* 5: 151-153 (1989); Corpet, *et al.*, *Nucleic Acids Research* 16: 10881-90 (1988); Huang, *et al.*, *Computer Applications in the Biosciences* 8: 155-65 (1992), and Pearson, *et al.*, *Methods in Molecular Biology* 24: 307-331 (1994). The BLAST family of programs which can be used for database similarity searches includes: BLASTN for nucleotide query sequences against nucleotide database sequences; BLASTX for nucleotide query sequences against protein database sequences; BLASTP for protein query sequences against protein database sequences; TBLASTN for protein query sequences against nucleotide database sequences; and TBLASTX for nucleotide query sequences against nucleotide database sequences. See, *Current Protocols in Molecular Biology*, Chapter 19, Ausubel, *et al.*, Eds., Greene Publishing and Wiley-Interscience, New York (1995).

GAP uses the algorithm of Needleman and Wunsch (*J. Mol. Biol.* 48: 443-453, 1970) to find the alignment of two complete sequences that maximizes the number of matches and minimizes the number of gaps. GAP considers all possible alignments and gap positions and creates the alignment with the largest number of matched bases and the fewest gaps. It allows for the provision of a gap creation penalty and a gap extension penalty in units of matched bases. GAP must make a profit of gap creation penalty number of matches for each gap it inserts. If a gap extension penalty greater than zero is chosen, GAP must, in addition, make a profit for each gap inserted of the length of the gap times the gap extension penalty. Default gap creation penalty values and gap extension penalty values in Version 10 of the Wisconsin Genetics Software Package for protein sequences are 8 and 2, respectively. For nucleotide sequences the default gap creation penalty is 50 while the default gap extension penalty is 3. The gap creation and

- 20 -

gap extension penalties can be expressed as an integer selected from the group of integers consisting of from 0 to 200. Thus, for example, the gap creation and gap extension penalties can be 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65 or greater.

5 GAP presents one member of the family of best alignments. There may be many members of this family, but no other member has a better quality. GAP displays four figures of merit for alignments: Quality, Ratio, Identity, and Similarity. The Quality is the metric maximized in order to align the sequences. Ratio is the quality divided by the number of bases in the shorter segment. Percent Identity is the percent of the symbols
10 that actually match. Percent Similarity is the percent of the symbols that are similar. Symbols that are across from gaps are ignored. A similarity is scored when the scoring matrix value for a pair of symbols is greater than or equal to 0.50, the similarity threshold. The scoring matrix used in Version 10 of the Wisconsin Genetics Software Package is BLOSUM62 (*see* Henikoff & Henikoff (1989) *Proc. Natl. Acad. Sci. USA*
15 89:10915).

 Unless otherwise stated, sequence identity/similarity values provided herein refer to the value obtained using the BLAST 2.0 suite of programs using default parameters (Altschul *et al.*, *Nucleic Acids Res.* 25:3389-3402, 1997; Altschul *et al.*, *J. Mol. Bio.* 215: 403-410, 1990) or to the value obtained using the GAP program using default
20 parameters (see the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, Wisconsin, USA).

 As those of ordinary skill in the art will understand, BLAST searches assume that proteins can be modeled as random sequences. However, many real proteins comprise regions of nonrandom sequences which may be homopolymeric tracts, short-period
25 repeats, or regions enriched in one or more amino acids. Such low-complexity regions may be aligned between unrelated proteins even though other regions of the protein are entirely dissimilar. A number of low-complexity filter programs can be employed to reduce such low-complexity alignments. For example, the SEG (Wooten and Federhen, *Comput. Chem.*, 17:149-163 (1993)) and XNU (Claverie and States, *Comput. Chem.*,
30 17:191-201 (1993)) low-complexity filters can be employed alone or in combination.

(c) As used herein, "sequence identity" or "identity" in the context of two nucleic acid or polypeptide sequences includes reference to the residues in the two sequences which are the same when aligned for maximum correspondence over a

- 21 -

specified comparison window. When percentage of sequence identity is used in reference to proteins it is recognized that residue positions which are not identical often differ by conservative amino acid substitutions, where amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g. charge or hydrophobicity) and therefore do not change the functional properties of the molecule. Where sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the substitution. Sequences which differ by such conservative substitutions are said to have "sequence similarity" or "similarity". Means for making this adjustment are well-known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated, e.g., according to the algorithm of Meyers and Miller, *Computer Applic. Biol. Sci.*, 4: 11-17 (1988) e.g., as implemented in the program PC/GENE (Intelligenetics, Mountain View, California, USA).

(d) As used herein, "percentage of sequence identity" means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

(e) (i) The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 70% sequence identity, preferably at least 80%, more preferably at least 90% and most preferably at least 95%, compared to a reference sequence using one of the alignment programs described using standard parameters. One of skill will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide

sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 60%, more preferably at least 70%, 80%, 90%, and most preferably at least 95%.

- 5 Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other under stringent conditions. However, nucleic acids which do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical. This may occur, *e.g.*, when a copy of a nucleic acid is created using the maximum codon
10 degeneracy permitted by the genetic code. One indication that two nucleic acid sequences are substantially identical is that the polypeptide which the first nucleic acid encodes is immunologically cross reactive with the polypeptide encoded by the second nucleic acid.

- (e) (ii) The terms "substantial identity" in the context of a peptide indicates that a peptide comprises a sequence with at least 70% sequence identity to a
15 reference sequence, preferably 80%, more preferably 85%, most preferably at least 90% or 95% sequence identity to the reference sequence over a specified comparison window. Preferably, optimal alignment is conducted using the homology alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48: 443 (1970). An indication that two peptide sequences are substantially identical is that one peptide is immunologically reactive with
20 antibodies raised against the second peptide. Thus, a peptide is substantially identical to a second peptide, for example, where the two peptides differ only by a conservative substitution. Peptides which are "substantially similar" share sequences as noted above except that residue positions which are not identical may differ by conservative amino acid changes.

25

DETAILED DESCRIPTION OF THE INVENTION

Overview

- The present invention provides, *inter alia*, compositions and methods for modulating (i.e., increasing or decreasing) the level of polypeptides of the present
30 invention in plants. In particular, the polypeptides of the present invention can be expressed at developmental stages, in tissues, and/or in quantities which are uncharacteristic of non-recombinantly engineered plants. Thus, the present invention

provides utility in such exemplary applications as the control of homologous recombination efficiency or transformation efficiency in plants.

The present invention also provides isolated nucleic acid comprising polynucleotides of sufficient length and complementarity to a gene of the present invention to use as probes or amplification primers in the detection, quantitation, or isolation of gene transcripts. For example, isolated nucleic acids of the present invention can be used as probes in detecting deficiencies in the level of mRNA in screenings for desired transgenic plants, for detecting mutations in the gene (e.g., substitutions, deletions, or additions), for monitoring upregulation of expression or changes in enzyme activity in screening assays of compounds, for detection of any number of allelic variants (polymorphisms) of the gene, or for use as molecular markers in plant breeding programs. The isolated nucleic acids of the present invention can also be used for recombinant expression of their encoded polypeptides, or for use as immunogens in the preparation and/or screening of antibodies. The isolated nucleic acids of the present invention can also be employed for use in sense or antisense suppression of one or more genes of the present invention in a host cell, tissue, or plant. Attachment of chemical agents which bind, intercalate, cleave and/or crosslink to the isolated nucleic acids of the present invention can also be used to modulate transcription or translation.

The present invention also provides isolated proteins comprising a polypeptide of the present invention (e.g., preproenzyme, proenzyme, or enzymes). The present invention also provides proteins comprising at least one epitope from a polypeptide of the present invention. The proteins of the present invention can be employed in assays for enzyme agonists or antagonists of enzyme function, or for use as immunogens or antigens to obtain antibodies specifically immunoreactive with a protein of the present invention. Such antibodies can be used in assays for expression levels, for identifying and/or isolating nucleic acids of the present invention from expression libraries, or for purification of polypeptides of the present invention.

The isolated nucleic acids and proteins of the present invention can be used over a broad range of plant types, particularly monocots such as the species of the Family *Graminae* including *Sorghum bicolor* and *Zea mays*. The isolated nucleic acid and proteins of the present invention can also be used in species from the genera: *Cucurbita*, *Rosa*, *Vitis*, *Juglans*, *Fragaria*, *Lotus*, *Medicago*, *Onobrychis*, *Trifolium*, *Trigonella*, *Vigna*, *Citrus*, *Linum*, *Geranium*, *Manihot*, *Daucus*, *Arabidopsis*, *Brassica*, *Raphanus*,

- 24 -

Sinapis, Atropa, Capsicum, Datura, Hyoscyamus, Lycopersicon, Nicotiana, Solanum, Petunia, Digitalis, Majorana, Cichorium, Helianthus, Lactuca, Bromus, Asparagus, Antirrhinum, Heterocallis, Nemesis, Pelargonium, Panieum, Pennisetum, Ranunculus, Senecio, Salpiglossis, Cucumis, Browaalia, Glycine, Pisum, Phaseolus, Lolium, Oryza,
5 *Avena, Hordeum, Secale, and Triticum.*

Nucleic Acids

The present invention provides, *inter alia*, isolated nucleic acids of RNA, DNA, and analogs and/or chimeras thereof, comprising a polynucleotide of the present
10 invention.

A polynucleotide of the present invention is inclusive of:

(a) a polynucleotide encoding a polypeptide of SEQ ID NO: 2 and conservatively modified and polymorphic variants thereof, including exemplary polynucleotides of SEQ ID NO: 1; polynucleotide sequences of the invention also
15 include the maize Ku70 polynucleotide sequence as contained in a plasmid deposited with American Type Culture Collection (ATCC) and assigned Accession Number _____.

(b) a polynucleotide which is the product of amplification from a *Zea mays* nucleic acid library using primer pairs which selectively hybridize under stringent
20 conditions to loci within the polynucleotide of SEQ ID NO: 1, or the sequence as contained in the ATCC deposit assigned Accession Number _____, wherein the polynucleotide has substantial sequence identity to the polynucleotide of SEQ ID NO: 1; or the sequence as contained in the ATCC deposit assigned Accession Number _____.

(c) a polynucleotide which selectively hybridizes to a polynucleotide of (a)
25 or (b);

(d) a polynucleotide having a specified sequence identity with polynucleotides of (a), (b), or (c);

(e) a polynucleotide encoding a protein having a specified number of contiguous amino acids from a prototype polypeptide, wherein the protein is specifically
30 recognized by antisera elicited by presentation of the protein and wherein the protein does not detectably immunoreact to antisera which has been fully immunosorbed with the protein;

- 25 -

(f) complementary sequences of polynucleotides of (a), (b), (c), (d), or (e); and

(g) a polynucleotide comprising at least a specific number of contiguous nucleotides from a polynucleotide of (a), (b), (c), (d), (e), or (f).

The polynucleotide of SEQ ID NO: 1 is contained in a plasmid deposited with American Type Culture Collection (ATCC) on August 26, 1999 and assigned Accession Number _____. American Type Culture Collection is located at 10801 University Blvd., Manassas, VA 20110-2209.

The ATCC deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. The deposit is provided as a convenience to those of skill in the art and is not an admission that a deposit is required under 35 U.S.C. Section 112. The deposited sequences, as well as the polypeptides encoded by the sequences, are incorporated herein by reference and control in the event of any conflict, such as a sequencing error, with the description in this application.

A. Polynucleotides Encoding A Polypeptide of the Present Invention or Conservatively Modified or Polymorphic Variants Thereof

As indicated in (a), *supra*, the present invention provides isolated nucleic acids comprising a polynucleotide of the present invention, wherein the polynucleotide encodes a polypeptide of the present invention, or conservatively modified or polymorphic variants thereof. Those of skill in the art will recognize that the degeneracy of the genetic code allows for a plurality of polynucleotides to encode for the identical amino acid sequence. Such "silent variations" can be used, for example, to selectively hybridize and detect allelic variants of polynucleotides of the present invention. Accordingly, the present invention includes polynucleotides of SEQ ID NO: 1, and the sequence as contained in the ATCC deposit assigned Accession Number _____, and silent variations of polynucleotides encoding a polypeptide of SEQ ID NO: 2. The present invention further provides isolated nucleic acids comprising polynucleotides encoding conservatively modified variants of a polypeptide of SEQ ID NO: 2. Conservatively modified variants can be used to generate or select antibodies immunoreactive to the non-variant polypeptide. Additionally, the present invention

- 26 -

further provides isolated nucleic acids comprising polynucleotides encoding one or more polymorphic (allelic) variants of polypeptides/polynucleotides. Polymorphic variants are frequently used to follow segregation of chromosomal regions in, for example, marker assisted selection methods for crop improvement.

5

B. Polynucleotides Amplified from a Zea mays Nucleic Acid Library

As indicated in (b), *supra*, the present invention provides an isolated nucleic acid comprising a polynucleotide of the present invention, wherein the polynucleotides are amplified from a *Zea mays* nucleic acid library. *Zea mays* lines B73, PHRE1, A632,
10 BMS-P2#10, W23, and Mo17 are known and publicly available. Other publicly known and available maize lines can be obtained from the Maize Genetics Cooperation (Urbana, IL). The nucleic acid library may be a cDNA library, a genomic library, or a library generally constructed from nuclear transcripts at any stage of intron processing. cDNA
15 libraries can be normalized to increase the representation of relatively rare cDNAs. In optional embodiments, the cDNA library is constructed using a full-length cDNA synthesis method. Examples of such methods include Oligo-Capping (Maruyama, K. and Sugano, S. *Gene* 138: 171-174, 1994), Biotinylated CAP Trapper (Carninci, P., Kvan, C., *et al.* *Genomics* 37: 327-336, 1996), and CAP Retention Procedure (Edery, E., Chu, L.L., *et al.* *Molecular and Cellular Biology* 15: 3363-3371, 1995). cDNA
20 synthesis is often catalyzed at 50-55°C to prevent formation of RNA secondary structure. Examples of reverse transcriptases that are relatively stable at these temperatures are SuperScript II Reverse Transcriptase (Life Technologies, Inc.), AMV Reverse Transcriptase (Boehringer Mannheim) and RetroAmp Reverse Transcriptase (Epicentre). Rapidly growing tissues, or rapidly dividing cells are preferably used as
25 mRNA sources.

The present invention also provides subsequences of the polynucleotides of the present invention. A variety of subsequences can be obtained using primers which selectively hybridize under stringent conditions to at least two sites within a polynucleotide of the present invention, or to two sites within the nucleic acid which
30 flank and comprise a polynucleotide of the present invention, or to a site within a polynucleotide of the present invention and a site within the nucleic acid which comprises it. Primers are chosen to selectively hybridize, under stringent hybridization conditions, to a polynucleotide of the present invention. Generally, the primers are

- 27 -

complementary to a subsequence of the target nucleic acid which they amplify. As those skilled in the art will appreciate, the sites to which the primer pairs will selectively hybridize are chosen such that a single contiguous nucleic acid can be formed under the desired amplification conditions. In optional embodiments, the primers will be

5 constructed so that they selectively hybridize under stringent conditions to a sequence (or its complement) within the target nucleic acid which comprises the codon encoding the carboxy or amino terminal amino acid residue (i.e., the 3' terminal coding region and 5' terminal coding region, respectively) of the polynucleotides of the present invention. Optionally within these embodiments, the primers will be constructed to selectively

10 hybridize entirely within the coding region of the target polynucleotide of the present invention such that the product of amplification of a cDNA target will consist of the coding region of that cDNA. The primer length in nucleotides is selected from the group of integers consisting of from at least 15 to 50. Thus, the primers can be at least 15, 18, 20, 25, 30, 40, or 50 nucleotides in length. Those of skill will recognize that a

15 lengthened primer sequence can be employed to increase specificity of binding (i.e., annealing) to a target sequence. A non-annealing sequence at the 5' end of a primer (a "tail") can be added, for example, to introduce a cloning site at the terminal ends of the amplicon.

The amplification products can be translated using expression systems well

20 known to those of skill in the art and as discussed, *infra*. The resulting translation products can be confirmed as polypeptides of the present invention by, for example, assaying for the appropriate catalytic activity (e.g., specific activity and/or substrate specificity), or verifying the presence of one or more linear epitopes which are specific to a polypeptide of the present invention. Methods for protein synthesis from PCR

25 derived templates are known in the art and available commercially. See, e.g., Amersham Life Sciences, Inc, Catalog '97, p.354.

Methods for obtaining 5' and/or 3' ends of a vector insert are well known in the art. See, e.g., RACE (Rapid Amplification of Complementary Ends) as described in Frohman, M. A., in PCR Protocols: A Guide to Methods and Applications, M. A. Innis,

30 D. H. Gelfand, J. J. Sninsky, T. J. White, Eds. (Academic Press, Inc., San Diego, 1990), pp. 28-38.); see also, U.S. Pat. No. 5,470,722, and *Current Protocols in Molecular Biology*, Unit 15.6, Ausubel, *et al.*, Eds., Greene Publishing and Wiley-Interscience, New York (1995); Frohman and Martin, *Techniques* 1:165 (1989).

C. Polynucleotides Which Selectively Hybridize to a Polynucleotide of (A) or (B)

As indicated in (c), *supra*, the present invention provides isolated nucleic acids comprising polynucleotides of the present invention, wherein the polynucleotides selectively hybridize, under selective hybridization conditions, to a polynucleotide of paragraphs (A) or (B) as discussed, *supra*. Thus, the polynucleotides of this embodiment can be used for isolating, detecting, and/or quantifying nucleic acids comprising the polynucleotides of (A) or (B). For example, polynucleotides of the present invention can be used to identify, isolate, or amplify partial or full-length clones in a deposited library. In some embodiments, the polynucleotides are genomic or cDNA sequences isolated or otherwise complementary to a cDNA from a dicot or monocot nucleic acid library. Exemplary species of monocots and dicots include, but are not limited to: corn, canola, soybean, cotton, wheat, sorghum, sunflower, oats, sugar cane, millet, barley, and rice. Preferably, the cDNA library comprises at least 80% full-length sequences, preferably at least 85% or 90% full-length sequences, and more preferably at least 95% full-length sequences. The cDNA libraries can be normalized to increase the representation of rare sequences. Low stringency hybridization conditions are typically, but not exclusively, employed with sequences having a reduced sequence identity relative to complementary sequences. Moderate and high stringency conditions can optionally be employed for sequences of greater identity. Low stringency conditions allow selective hybridization of sequences having about 70% sequence identity and can be employed to identify orthologous or paralogous sequences.

D. Polynucleotides Having a Specific Sequence Identity with the Polynucleotides of (A), (B) or (C)

As indicated in (d), *supra*, the present invention provides isolated nucleic acids comprising polynucleotides of the present invention, wherein the polynucleotides have a specified identity at the nucleotide level to a polynucleotide as disclosed above in paragraphs (A), (B), or (C). The percentage of identity to a reference sequence is at least 60% and, rounded upwards to the nearest integer, can be expressed as an integer selected from the group of integers consisting of from 60 to 99. Thus, for example, the percentage of identity to a reference sequence can be at least 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%.

Optionally, the polynucleotides of this embodiment will share an epitope with a polypeptide encoded by the polynucleotides of (A), (B), or (C). Thus, these polynucleotides encode a first polypeptide which elicits production of antisera comprising antibodies which are specifically reactive to a second polypeptide encoded by a polynucleotide of (A), (B), or (C). However, the first polypeptide does not bind to antisera raised against itself when the antisera has been fully immunosorbed with the first polypeptide. Hence, the polynucleotides of this embodiment can be used to generate antibodies for use in, for example, the screening of expression libraries for nucleic acids comprising polynucleotides of (A), (B), or (C), or for purification of, or in immunoassays for, polypeptides encoded by the polynucleotides of (A), (B), or (C). The polynucleotides of this embodiment embrace nucleic acid sequences which can be employed for selective hybridization to a polynucleotide encoding a polypeptide of the present invention.

Screening polypeptides for specific binding to antisera can be conveniently achieved using peptide display libraries. This method involves the screening of large collections of peptides for individual members having the desired function or structure. Antibody screening of peptide display libraries is well known in the art. The displayed peptide sequences can be from 3 to 5000 or more amino acids in length, frequently from 5-100 amino acids long, and often from about 8 to 15 amino acids long. In addition to direct chemical synthetic methods for generating peptide libraries, several recombinant DNA methods have been described. One type involves the display of a peptide sequence on the surface of a bacteriophage or cell. Each bacteriophage or cell contains the nucleotide sequence encoding the particular displayed peptide sequence. Such methods are described in PCT patent publication Nos. 91/17271, 91/18980, 91/19818, and 93/08278. Other systems for generating libraries of peptides have aspects of both *in vitro* chemical synthesis and recombinant methods. See, PCT Patent publication Nos. 92/05258, 92/14843, and 96/19256. See also, U.S. Patent Nos. 5,658,754; and 5,643,768. Peptide display libraries, vectors, and screening kits are commercially available from such suppliers as Invitrogen (Carlsbad, CA).

E. Polynucleotides Encoding a Protein Having a Subsequence from a Prototype Polypeptide and is Cross-Reactive to the Prototype Polypeptide

As indicated in (e), *supra*, the present invention provides isolated nucleic acids comprising polynucleotides of the present invention, wherein the polynucleotides encode
5 a protein having a subsequence of contiguous amino acids from a prototype polypeptide of the present invention such as are provided in (a), *supra*. The length of contiguous amino acids from the prototype polypeptide is selected from the group of integers consisting of from at least 10 to the number of amino acids within the prototype sequence. Thus, for example, the polynucleotide can encode a polypeptide having a
10 subsequence having at least 10, 15, 20, 25, 30, 35, 40, 45, or 50, contiguous amino acids from the prototype polypeptide. Further, the number of such subsequences encoded by a polynucleotide of the instant embodiment can be any integer selected from the group consisting of from 1 to 20, such as 2, 3, 4, or 5. The subsequences can be separated by any integer of nucleotides from 1 to the number of nucleotides in the
15 sequence such as at least 5, 10, 15, 25, 50, 100, or 200 nucleotides.

The proteins encoded by polynucleotides of this embodiment, when presented as an immunogen, elicit the production of polyclonal antibodies which specifically bind to a prototype polypeptide such as but not limited to, a polypeptide encoded by the polynucleotide of (a) or (b), *supra*. Generally, however, a protein encoded by a
20 polynucleotide of this embodiment does not bind to antisera raised against the prototype polypeptide when the antisera has been fully immunosorbed with the prototype polypeptide. Methods of making and assaying for antibody binding specificity/affinity are well known in the art. Exemplary immunoassay formats include ELISA, competitive immunoassays, radioimmunoassays, Western blots, indirect immunofluorescent assays
25 and the like.

In a preferred assay method, fully immunosorbed and pooled antisera which is elicited to the prototype polypeptide can be used in a competitive binding assay to test the protein. The concentration of the prototype polypeptide required to inhibit 50% of the binding of the antisera to the prototype polypeptide is determined. If the amount of
30 the protein required to inhibit binding is less than twice the amount of the prototype protein, then the protein is said to specifically bind to the antisera elicited to the immunogen. Accordingly, the proteins of the present invention embrace allelic variants,

conservatively modified variants, and minor recombinant modifications to a prototype polypeptide.

A polynucleotide of the present invention optionally encodes a protein having a molecular weight as the non-glycosylated protein within 20% of the molecular weight of the full-length non-glycosylated polypeptides of the present invention. Molecular weight can be readily determined by SDS-PAGE under reducing conditions. Preferably, the molecular weight is within 15% of a full length polypeptide of the present invention, more preferably within 10% or 5%, and most preferably within 3%, 2%, or 1% of a full length polypeptide of the present invention. Molecular weight determination of a protein can be conveniently performed by SDS-PAGE under denaturing conditions.

Optionally, the polynucleotides of this embodiment will encode a protein having a specific activity at least 50%, 60%, 80%, or 90% of the native, endogenous (i.e., non-isolated), full-length polypeptide of the present invention. Further, the proteins encoded by polynucleotides of this embodiment will optionally have a substantially similar affinity constant (K_m) and/or catalytic activity (i.e., the microscopic rate constant, k_{cat}) as the native endogenous, full-length protein. Those of skill in the art will recognize that k_{cat}/K_m value determines the specificity for competing substrates and is often referred to as the specificity constant. Proteins of this embodiment can have a k_{cat}/K_m value at least 10% of a non-isolated full-length polypeptide of the present invention as determined using the endogenous substrate of that polypeptide. Optionally, the k_{cat}/K_m value will be at least 20%, 30%, 40%, 50%, and most preferably at least 60%, 70%, 80%, 90%, or 95% the k_{cat}/K_m value of the non-isolated, full-length polypeptide of the present invention.

Determination of k_{cat} , K_m , and k_{cat}/K_m can be determined by any number of means well known to those of skill in the art. For example, the initial rates (i.e., the first 5% or less of the reaction) can be determined using rapid mixing and sampling techniques (e.g., continuous-flow, stopped-flow, or rapid quenching techniques), flash photolysis, or relaxation methods (e.g., temperature jumps) in conjunction with such exemplary methods of measuring as spectrophotometry, spectrofluorimetry, nuclear magnetic resonance, or radioactive procedures. Kinetic values are conveniently obtained using a Lineweaver-Burk or Eadie-Hofstee plot.

F. Polynucleotides Complementary to the Polynucleotides of (A)-(E)

As indicated in (f), *supra*, the present invention provides isolated nucleic acids comprising polynucleotides complementary to the polynucleotides of paragraphs A-E, above. As those of skill in the art will recognize, complementary sequences base-pair
5 throughout the entirety of their length with the polynucleotides of (A)-(E) (i.e., have 100% sequence identity over their entire length). Complementary bases associate through hydrogen bonding in double stranded nucleic acids. For example, the following base pairs are complementary: guanine and cytosine; adenine and thymine; and adenine and uracil.

10

G. Polynucleotides Which are Subsequences of the Polynucleotides of (A)-(F)

As indicated in (g), *supra*, the present invention provides isolated nucleic acids comprising polynucleotides which comprise at least 15 contiguous bases from the polynucleotides of (A) through (F) as discussed above. The length of the polynucleotide
15 is given as an integer selected from the group consisting of from at least 15 to the length of the nucleic acid sequence from which the polynucleotide is a subsequence of. Thus, for example, polynucleotides of the present invention are inclusive of polynucleotides comprising at least 15, 20, 25, 30, 40, 50, 60, 75, or 100 contiguous nucleotides in length from the polynucleotides of (A)-(F). Optionally, the number of such
20 subsequences encoded by a polynucleotide of the instant embodiment can be any integer selected from the group consisting of from 1 to 20, such as 2, 3, 4, or 5. The subsequences can be separated by any integer of nucleotides from 1 to the number of nucleotides in the sequence such as at least 5, 10, 15, 25, 50, 100, or 200 nucleotides.

The subsequences of the present invention can comprise structural characteristics
25 of the sequence from which it is derived. Alternatively, the subsequences can lack certain structural characteristics of the larger sequence from which it is derived. For example, a subsequence from a polynucleotide encoding a polypeptide having at least one linear epitope in common with a prototype polypeptide sequence as provided in (a), *supra*, may encode an epitope in common with the prototype sequence. Alternatively,
30 the subsequence may not encode an epitope in common with the prototype sequence but can be used to isolate the larger sequence by, for example, nucleic acid hybridization with the sequence from which it's derived. Subsequences can be used to modulate or detect gene expression by introducing into the subsequences compounds which bind,

intercalate, cleave and/or crosslink to nucleic acids. Exemplary compounds include acridine, psoralen, phenanthroline, naphthoquinone, daunomycin or chloroethylaminoaryl conjugates.

5 Construction of Nucleic Acids

The isolated nucleic acids of the present invention can be made using (a) standard recombinant methods, (b) synthetic techniques, or combinations thereof. In some embodiments, the polynucleotides of the present invention will be cloned, amplified, or otherwise constructed from a monocot. In preferred embodiments the monocot is *Zea mays*.

The nucleic acids may conveniently comprise sequences in addition to a polynucleotide of the present invention. For example, a multi-cloning site comprising one or more endonuclease restriction sites may be inserted into the nucleic acid to aid in isolation of the polynucleotide. Also, translatable sequences may be inserted to aid in the isolation of the translated polynucleotide of the present invention. For example, a hexa-histidine marker sequence provides a convenient means to purify the proteins of the present invention. A polynucleotide of the present invention can be attached to a vector, adapter, or linker for cloning and/or expression of a polynucleotide of the present invention. Additional sequences may be added to such cloning and/or expression sequences to optimize their function in cloning and/or expression, to aid in isolation of the polynucleotide, or to improve the introduction of the polynucleotide into a cell. Typically, the length of a nucleic acid of the present invention less the length of its polynucleotide of the present invention is less than 20 kilobase pairs, often less than 15 kb, and frequently less than 10 kb. Use of cloning vectors, expression vectors, adapters, and linkers is well known and extensively described in the art. For a description of various nucleic acids see, for example, Stratagene Cloning Systems, Catalogs 1995, 1996, 1997 (La Jolla, CA); and, Amersham Life Sciences, Inc, Catalog '97 (Arlington Heights, IL).

30 *A. Recombinant Methods for Constructing Nucleic Acids*

The isolated nucleic acid compositions of this invention, such as RNA, cDNA, genomic DNA, or a hybrid thereof, can be obtained from plant biological sources using any number of cloning methodologies known to those of skill in the art. In some

embodiments, oligonucleotide probes which selectively hybridize, under stringent conditions, to the polynucleotides of the present invention are used to identify the desired sequence in a cDNA or genomic DNA library. While isolation of RNA, and construction of cDNA and genomic libraries is well known to those of ordinary skill in the art, the following highlights some of the methods employed.

A1. mRNA Isolation and Purification

Total RNA from plant cells comprises such nucleic acids as mitochondrial RNA, chloroplastic RNA, rRNA, tRNA, hnRNA and mRNA. Total RNA preparation typically involves lysis of cells and removal of proteins, followed by precipitation of nucleic acids. Extraction of total RNA from plant cells can be accomplished by a variety of means. Frequently, extraction buffers include a strong detergent such as SDS and an organic denaturant such as guanidinium isothiocyanate, guanidine hydrochloride or phenol. Following total RNA isolation, poly(A)⁺ mRNA is typically purified from the remainder RNA using oligo(dT) cellulose. Exemplary total RNA and mRNA isolation protocols are described in *Plant Molecular Biology: A Laboratory Manual*, Clark, Ed., Springer-Verlag, Berlin (1997); and, *Current Protocols in Molecular Biology*, Ausubel, *et al.*, Eds., Greene Publishing and Wiley-Interscience, New York (1995). Total RNA and mRNA isolation kits are commercially available from vendors such as Stratagene (La Jolla, CA), Clontech (Palo Alto, CA), Pharmacia (Piscataway, NJ), and 5'-3' (Paoli, PA). See also, U.S. Patent Nos. 5,614,391; and, 5,459,253. The mRNA can be fractionated into populations with size ranges of about 0.5, 1.0, 1.5, 2.0, 2.5 or 3.0 kb. The cDNA synthesized for each of these fractions can be size selected to the same size range as its mRNA prior to vector insertion. This method helps eliminate truncated cDNA formed by incompletely reverse transcribed mRNA.

A2. Construction of a cDNA Library

Construction of a cDNA library generally entails five steps. First, first strand cDNA synthesis is initiated from a poly(A)⁺ mRNA template using a poly(dT) primer or random hexanucleotides. Second, the resultant RNA-DNA hybrid is converted into double stranded cDNA, typically by a combination of RNase H and DNA polymerase I (or Klenow fragment). Third, the termini of the double stranded cDNA are ligated to adaptors. Ligation of the adaptors will produce cohesive ends for cloning. Fourth, size

selection of the double stranded cDNA eliminates excess adaptors and primer fragments, and eliminates partial cDNA molecules due to degradation of mRNAs or the failure of reverse transcriptase to synthesize complete first strands. Fifth, the cDNAs are ligated into cloning vectors and packaged. cDNA synthesis protocols are well known to the skilled artisan and are described in such standard references as: *Plant Molecular Biology: A Laboratory Manual*, Clark, Ed., Springer-Verlag, Berlin (1997); and, *Current Protocols in Molecular Biology*, Ausubel, *et al.*, Eds., Greene Publishing and Wiley-Interscience, New York (1995). cDNA synthesis kits are available from a variety of commercial vendors such as Stratagene or Pharmacia.

10 A number of cDNA synthesis protocols have been described which provide substantially pure full-length cDNA libraries. Substantially pure full-length cDNA libraries are constructed to comprise at least 90%, and more preferably at least 93% or 95% full-length inserts amongst clones containing inserts. The length of insert in such libraries can be from 0 to 8, 9, 10, 11, 12, 13, or more kilobase pairs. Vectors to
15 accommodate inserts of these sizes are known in the art and available commercially. See, e.g., Stratagene's lambda ZAP Express (cDNA cloning vector with 0 to 12 kb cloning capacity).

 An exemplary method of constructing a greater than 95% pure full-length cDNA library is described by Carninci *et al.*, *Genomics*, 37:327-336 (1996). In that protocol,
20 the cap-structure of eukaryotic mRNA is chemically labeled with biotin. By using streptavidin-coated magnetic beads, only the full-length first-strand cDNA/mRNA hybrids are selectively recovered after RNase I treatment. The method provides a high yield library with an unbiased representation of the starting mRNA population. Other methods for producing full-length libraries are known in the art. See, e.g., Edery *et al.*,
25 *Mol. Cell Biol.*, 15(6):3363-3371 (1995); and, PCT Application WO 96/34981.

A3. Normalized or Subtracted cDNA Libraries

 A non-normalized cDNA library represents the mRNA population of the tissue it was made from. Since unique clones are out-numbered by clones derived from highly
30 expressed genes their isolation can be laborious. Normalization of a cDNA library is the process of creating a library in which each clone is more equally represented.

 A number of approaches to normalize cDNA libraries are known in the art. One approach is based on hybridization to genomic DNA. The frequency of each hybridized

cDNA in the resulting normalized library would be proportional to that of each corresponding gene in the genomic DNA. Another approach is based on kinetics. If cDNA reannealing follows second-order kinetics, rarer species anneal less rapidly and the remaining single-stranded fraction of cDNA becomes progressively more normalized during the course of the hybridization. Specific loss of any species of cDNA, regardless of its abundance, does not occur at any Cot value. Construction of normalized libraries is described in Ko, *Nucl. Acids. Res.*, 18(19):5705-5711 (1990); Patanjali *et al.*, *Proc. Natl. Acad. U.S.A.*, 88:1943-1947 (1991); U.S. Patents 5,482,685, and 5,637,685. In an exemplary method described by Soares *et al.*, normalization resulted in reduction of the abundance of clones from a range of four orders of magnitude to a narrow range of only 1 order of magnitude. *Proc. Natl. Acad. Sci. USA*, 91:9228-9232 (1994).

Subtracted cDNA libraries are another means to increase the proportion of less abundant cDNA species. In this procedure, cDNA prepared from one pool of mRNA is depleted of sequences present in a second pool of mRNA by hybridization. The cDNA:mRNA hybrids are removed and the remaining un-hybridized cDNA pool is enriched for sequences unique to that pool. See, Foote *et al.* in, *Plant Molecular Biology: A Laboratory Manual*, Clark, Ed., Springer-Verlag, Berlin (1997); Kho and Zarbl, *Technique*, 3(2):58-63 (1991); Sive and St. John, *Nucl. Acids Res.*, 16(22):10937 (1988); *Current Protocols in Molecular Biology*, Ausubel, *et al.*, Eds., Greene Publishing and Wiley-Interscience, New York (1995); and, Swaroop *et al.*, *Nucl. Acids Res.*, 19(8):1954 (1991). cDNA subtraction kits are commercially available. See, e.g., PCR-Select (Clontech).

A4. Construction of a Genomic Library

To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation, e.g. using restriction endonucleases, and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. Methodologies to accomplish these ends, and sequencing methods to verify the sequence of nucleic acids are well known in the art. Examples of appropriate molecular biological techniques and instructions sufficient to direct persons of skill through many construction, cloning, and screening methodologies are found in Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Vols. 1-3 (1989), *Methods in Enzymology*, Vol. 152: *Guide to Molecular Cloning*

Techniques, Berger and Kimmel, Eds., San Diego: Academic Press, Inc. (1987), *Current Protocols in Molecular Biology*, Ausubel, *et al.*, Eds., Greene Publishing and Wiley-Interscience, New York (1995); *Plant Molecular Biology: A Laboratory Manual*, Clark, Ed., Springer-Verlag, Berlin (1997). Kits for construction of genomic libraries are also
5 commercially available.

A5. Nucleic Acid Screening and Isolation Methods

The cDNA or genomic library can be screened using a probe based upon the sequence of a polynucleotide of the present invention such as those disclosed herein.

10 Probes may be used to hybridize with genomic DNA or cDNA sequences to isolate homologous genes in the same or different plant species. Those of skill in the art will appreciate that various degrees of stringency of hybridization can be employed in the assay; and either the hybridization or the wash medium can be stringent. As the conditions for hybridization become more stringent, there must be a greater degree of
15 complementarity between the probe and the target for duplex formation to occur. The degree of stringency can be controlled by temperature, ionic strength, pH and the presence of a partially denaturing solvent such as formamide. For example, the stringency of hybridization is conveniently varied by changing the polarity of the reactant solution through manipulation of the concentration of formamide within the range of 0%
20 to 50%. The degree of complementarity (sequence identity) required for detectable binding will vary in accordance with the stringency of the hybridization medium and/or wash medium. The degree of complementarity will optimally be 100 percent; however, it should be understood that minor sequence variations in the probes and primers may be compensated for by reducing the stringency of the hybridization and/or wash medium.

25 The nucleic acids of interest can also be amplified from nucleic acid samples using amplification techniques. For instance, polymerase chain reaction (PCR) technology can be used to amplify the sequences of polynucleotides of the present invention and related genes directly from genomic DNA or cDNA libraries. PCR and other *in vitro* amplification methods may also be useful, for example, to clone nucleic
30 acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes. Examples of techniques sufficient to direct persons of skill through *in vitro* amplification methods are found in Berger, Sambrook, and

Ausubel, as well as Mullis *et al.*, U.S. Patent No. 4,683,202 (1987); and, *PCR Protocols A Guide to Methods and Applications*, Innis *et al.*, Eds., Academic Press Inc., San Diego, CA (1990). Commercially available kits for genomic PCR amplification are known in the art. See, e.g., Advantage-GC Genomic PCR Kit (Clontech). The T4 gene
5 32 protein (Boehringer Mannheim) can be used to improve yield of long PCR products.

PCR-based screening methods have also been described. Wilfinger *et al.* describe a PCR-based method in which the longest cDNA is identified in the first step so that incomplete clones can be eliminated from study. *BioTechniques*, 22(3): 481-486 (1997). In that method, a primer pair is synthesized with one primer annealing to the 5'
10 end of the sense strand of the desired cDNA and the other primer to the vector. Clones are pooled to allow large-scale screening. By this procedure, the longest possible clone is identified amongst candidate clones. Further, the PCR product is used solely as a diagnostic for the presence of the desired cDNA and does not utilize the PCR product itself. Such methods are particularly effective in combination with a full-length cDNA
15 construction methodology, *supra*.

B. Synthetic Methods for Constructing Nucleic Acids

The isolated nucleic acids of the present invention can also be prepared by direct chemical synthesis by methods such as the phosphotriester method of Narang *et al.*,
20 *Meth. Enzymol.* 68: 90-99 (1979); the phosphodiester method of Brown *et al.*, *Meth. Enzymol.* 68: 109-151 (1979); the diethylphosphoramidite method of Beaucage *et al.*, *Tetra. Lett.* 22: 1859-1862 (1981); the solid phase phosphoramidite triester method described by Beaucage and Caruthers, *Tetra. Letts.* 22(20): 1859-1862 (1981), e.g., using an automated synthesizer, e.g., as described in Needham-VanDevanter *et al.*,
25 *Nucleic Acids Res.*, 12: 6159-6168 (1984); and, the solid support method of U.S. Patent No. 4,458,066. Chemical synthesis generally produces a single stranded oligonucleotide. This may be converted into double stranded DNA by hybridization with a complementary sequence, or by polymerization with a DNA polymerase using the single strand as a template. One of skill will recognize that while chemical synthesis of
30 DNA is limited to sequences of about 100 bases, longer sequences may be obtained by the ligation of shorter sequences.

Recombinant Expression Cassettes

The present invention further provides recombinant expression cassettes comprising a nucleic acid of the present invention. A nucleic acid sequence coding for the desired polynucleotide of the present invention, for example a cDNA or a genomic
5 sequence encoding a full length polypeptide of the present invention, can be used to construct a recombinant expression cassette which can be introduced into the desired host cell. A recombinant expression cassette will typically comprise a polynucleotide of the present invention operably linked to transcriptional initiation regulatory sequences which will direct the transcription of the polynucleotide in the intended host cell, such as tissues
10 of a transformed plant.

For example, plant expression vectors may include (1) a cloned plant gene under the transcriptional control of 5' and 3' regulatory sequences and (2) a dominant selectable marker. Such plant expression vectors may also contain, if desired, a promoter regulatory region (e.g., one conferring inducible or constitutive, environmentally- or
15 developmentally-regulated, or cell- or tissue-specific/selective expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

A plant promoter fragment can be employed which will direct expression of a polynucleotide of the present invention in all tissues of a regenerated plant. Such
20 promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, the ubiquitin 1 promoter, the Smas promoter, the cinnamyl alcohol
25 dehydrogenase promoter (U.S. Patent No. 5,683,439), the *Nos* promoter, the pEmu promoter, the rubisco promoter, the GRP1-8 promoter, and other transcription initiation regions from various plant genes known to those of skill.

Alternatively, the plant promoter can direct expression of a polynucleotide of the present invention in a specific tissue or may be otherwise under more precise
30 environmental or developmental control. Such promoters are referred to here as "inducible" promoters. Environmental conditions that may effect transcription by inducible promoters include pathogen attack, anaerobic conditions, or the presence of light. Examples of inducible promoters are the *Adh1* promoter which is inducible by

hypoxia or cold stress, the Hsp70 promoter which is inducible by heat stress, and the PPKK promoter which is inducible by light.

Examples of promoters under developmental control include promoters that initiate transcription only, or preferentially, in certain tissues, such as leaves, roots,
5 fruit, seeds, or flowers. An exemplary promoter is the anther specific promoter 5126 (U.S. Patent Nos. 5,689,049 and 5,689,051). The operation of a promoter may also vary depending on its location in the genome. Thus, an inducible promoter may become fully or partially constitutive in certain locations.

Both heterologous and non-heterologous (i.e., endogenous) promoters can be
10 employed to direct expression of the nucleic acids of the present invention. These promoters can also be used, for example, in recombinant expression cassettes to drive expression of antisense nucleic acids to reduce, increase, or alter concentration and/or composition of the proteins of the present invention in a desired tissue. Thus, in some embodiments, the nucleic acid construct will comprise a promoter functional in a plant
15 cell, such as in *Zea mays*, operably linked to a polynucleotide of the present invention. Promoters useful in these embodiments include the endogenous promoters driving expression of a polypeptide of the present invention.

In some embodiments, isolated nucleic acids which serve as promoter or enhancer elements can be introduced in the appropriate position (generally upstream) of a non-
20 heterologous form of a polynucleotide of the present invention so as to up or down regulate expression of a polynucleotide of the present invention. For example, endogenous promoters can be altered *in vivo* by mutation, deletion, and/or substitution (see, Kmiec, U.S. Patent 5,565,350; Zarling *et al.*, PCT/US93/03868), or isolated promoters can be introduced into a plant cell in the proper orientation and distance from
25 a gene of the present invention so as to control the expression of the gene. Gene expression can be modulated under conditions suitable for plant growth so as to alter the total concentration and/or alter the composition of the polypeptides of the present invention in plant cell. Thus, the present invention provides compositions, and methods for making, heterologous promoters and/or enhancers operably linked to a native,
30 endogenous (i.e., non-heterologous) form of a polynucleotide of the present invention.

Methods for identifying promoters with a particular expression pattern, in terms of, e.g., tissue type, cell type, stage of development, and/or environmental conditions, are well known in the art. See, e.g., *The Maize Handbook*, Chapters 114-115, Freeling

and Walbot, Eds., Springer, New York (1994); *Corn and Corn Improvement*, 3rd edition, Chapter 6, Sprague and Dudley, Eds., American Society of Agronomy, Madison, Wisconsin (1988). A typical step in promoter isolation methods is identification of gene products that are expressed with some degree of specificity in the target tissue. Amongst
5 the range of methodologies are: differential hybridization to cDNA libraries; subtractive hybridization; differential display; differential 2-D protein gel electrophoresis; DNA probe arrays; and isolation of proteins known to be expressed with some specificity in the target tissue. Such methods are well known to those of skill in the art. Commercially available products for identifying promoters are known in the art such as
10 Clontech's (Palo Alto, CA) Universal GenomeWalker Kit.

For the protein-based methods, it is helpful to obtain the amino acid sequence for at least a portion of the identified protein, and then to use the protein sequence as the basis for preparing a nucleic acid that can be used as a probe to identify either genomic DNA directly, or preferably, to identify a cDNA clone from a library prepared from the
15 target tissue. Once such a cDNA clone has been identified, that sequence can be used to identify the sequence at the 5' end of the transcript of the indicated gene. For differential hybridization, subtractive hybridization and differential display, the nucleic acid sequence identified as enriched in the target tissue is used to identify the sequence at the 5' end of the transcript of the indicated gene. Once such sequences are identified,
20 starting either from protein sequences or nucleic acid sequences, any of these sequences identified as being from the gene transcript can be used to screen a genomic library prepared from the target organism. Methods for identifying and confirming the transcriptional start site are well known in the art.

In the process of isolating promoters expressed under particular environmental
25 conditions or stresses, or in specific tissues, or at particular developmental stages, a number of genes are identified that are expressed under the desired circumstances, in the desired tissue, or at the desired stage. Further analysis will reveal expression of each particular gene in one or more other tissues of the plant. One can identify a promoter with activity in the desired tissue or condition but that do not have activity in any other
30 common tissue.

To identify the promoter sequence, the 5' portions of the clones described here are analyzed for sequences characteristic of promoter sequences. For instance, promoter sequence elements include the TATA box consensus sequence (TATAAT), which is

usually an AT-rich stretch of 5-10 bp located approximately 20 to 40 base pairs upstream of the transcription start site. Identification of the TATA box is well known in the art. For example, one way to predict the location of this element is to identify the transcription start site using standard RNA-mapping techniques such as primer extension, S1 analysis, and/or RNase protection. To confirm the presence of the AT-rich sequence, a structure-function analysis can be performed involving mutagenesis of the putative region and quantification of the mutation's effect on expression of a linked downstream reporter gene. See, e.g., *The Maize Handbook*, Chapter 114, Freeling and Walbot, Eds., Springer, New York, (1994).

10 In plants, further upstream from the TATA box, at positions -80 to -100, there is typically a promoter element (i.e., the CAAT box) with a series of adenines surrounding the trinucleotide G (or T) N G. J. Messing *et al.*, in *Genetic Engineering in Plants*, Kosage, Meredith and Hollaender, Eds., pp. 221-227 1983. In maize, there is no well conserved CAAT box but there are several short, conserved protein-binding motifs upstream of the TATA box. These include motifs for the trans-acting transcription factors involved in light regulation, anaerobic induction, hormonal regulation, or anthocyanin biosynthesis, as appropriate for each gene.

Once promoter and/or gene sequences are known, a region of suitable size is selected from the genomic DNA that is 5' to the transcriptional start, or the translational start site, and such sequences are then linked to a coding sequence. If the transcriptional start site is used as the point of fusion, any of a number of possible 5' untranslated regions can be used in between the transcriptional start site and the partial coding sequence. If the translational start site at the 3' end of the specific promoter is used, then it is linked directly to the methionine start codon of a coding sequence.

25 If polypeptide expression is desired, it is generally desirable to include a polyadenylation region at the 3'-end of a polynucleotide coding region. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA. The 3' end sequence to be added can be derived from, for example, the nopaline synthase or octopine synthase genes, or alternatively from another plant gene, or less preferably from any other eukaryotic gene.

30 An intron sequence can be added to the 5' untranslated region or the coding sequence of the partial coding sequence to increase the amount of the mature message that accumulates in the cytosol. Inclusion of a spliceable intron in the transcription unit

in both plant and animal expression constructs has been shown to increase gene expression at both the mRNA and protein levels up to 1000-fold. Buchman and Berg, *Mol. Cell Biol.* 8: 4395-4405 (1988); Callis *et al.*, *Genes Dev.* 1: 1183-1200 (1987). Such intron enhancement of gene expression is typically greatest when placed near the 5' end of the transcription unit. Use of maize introns Adh1-S intron 1, 2, and 6, the Bronze-1 intron are known in the art. See generally, *The Maize Handbook*, Chapter 116, Freeling and Walbot, Eds., Springer, New York (1994).

The vector comprising the sequences from a polynucleotide of the present invention will typically comprise a marker gene which confers a selectable phenotype on plant cells. Usually, the selectable marker gene will encode antibiotic resistance, with suitable genes including genes coding for resistance to the antibiotic spectinomycin (e.g., the *aada* gene), the streptomycin phosphotransferase (SPT) gene coding for streptomycin resistance, the neomycin phosphotransferase (NPTII) gene encoding kanamycin or geneticin resistance, the hygromycin phosphotransferase (HPT) gene coding for hygromycin resistance, genes coding for resistance to herbicides which act to inhibit the action of acetolactate synthase (ALS), in particular the sulfonyleurea-type herbicides (e.g., the acetolactate synthase (ALS) gene containing mutations leading to such resistance in particular the S4 and/or Hra mutations), genes coding for resistance to herbicides which act to inhibit action of glutamine synthase, such as phosphinothricin or basta (e.g., the *bar* gene), or other such genes known in the art. The *bar* gene encodes resistance to the herbicide basta, the *nptII* gene encodes resistance to the antibiotics kanamycin and geneticin, and the ALS gene encodes resistance to the herbicide chlorsulfuron.

Typical vectors useful for expression of genes in higher plants are well known in the art and include vectors derived from the tumor-inducing (Ti) plasmid of *Agrobacterium tumefaciens* described by Rogers *et al.*, *Meth. in Enzymol.*, 153:253-277 (1987). These vectors are plant integrating vectors in that on transformation, the vectors integrate a portion of vector DNA into the genome of the host plant. Exemplary *A. tumefaciens* vectors useful herein are plasmids pKYLX6 and pKYLX7 of Schardl *et al.*, *Gene*, 61:1-11 (1987) and Berger *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 86:8402-8406 (1989). Another useful vector herein is plasmid pBI101.2 that is available from Clontech Laboratories, Inc. (Palo Alto, CA).

A polynucleotide of the present invention can be expressed in either sense or anti-sense orientation as desired. It will be appreciated that control of gene expression in either sense or anti-sense orientation can have a direct impact on the observable plant characteristics. Antisense technology can be conveniently used to gene expression in
5 plants. To accomplish this, a nucleic acid segment from the desired gene is cloned and operably linked to a promoter such that the anti-sense strand of RNA will be transcribed. The construct is then transformed into plants and the antisense strand of RNA is produced. In plant cells, it has been shown that antisense RNA inhibits gene expression by preventing the accumulation of mRNA which encodes the enzyme of interest, see,
10 e.g., Sheehy *et al.*, *Proc. Nat'l. Acad. Sci. (USA)* 85: 8805-8809 (1988); and Hiatt *et al.*, U.S. Patent No. 4,801,340.

Another method of suppression is sense suppression. Introduction of nucleic acid configured in the sense orientation has been shown to be an effective means by which to block the transcription of target genes. For an example of the use of this method to
15 modulate expression of endogenous genes see, Napoli *et al.*, *The Plant Cell* 2: 279-289 (1990) and U.S. Patent No. 5,034,323.

Catalytic RNA molecules or ribozymes can also be used to inhibit expression of plant genes. It is possible to design ribozymes that specifically pair with virtually any target RNA and cleave the phosphodiester backbone at a specific location, thereby
20 functionally inactivating the target RNA. In carrying out this cleavage, the ribozyme is not itself altered, and is thus capable of recycling and cleaving other molecules, making it a true enzyme. The inclusion of ribozyme sequences within antisense RNAs confers RNA-cleaving activity upon them, thereby increasing the activity of the constructs. The design and use of target RNA-specific ribozymes is described in Haseloff *et al.*, *Nature*
25 334: 585-591 (1988).

A variety of cross-linking agents, alkylating agents and radical generating species as pendant groups on polynucleotides of the present invention can be used to bind, label, detect, and/or cleave nucleic acids. For example, Vlassov, V. V., *et al.*, *Nucleic Acids Res* (1986) 14:4065-4076, describe covalent bonding of a single-stranded DNA fragment
30 with alkylating derivatives of nucleotides complementary to target sequences. A report of similar work by the same group is that by Knorre, D. G., *et al.*, *Biochimie* (1985) 67:785-789. Iverson and Dervan also showed sequence-specific cleavage of single-stranded DNA mediated by incorporation of a modified nucleotide which was capable of

- 45 -

activating cleavage (*J Am Chem Soc* (1987) 109:1241-1243). Meyer, R. B., *et al.*, *J Am Chem Soc* (1989) 111:8517-8519, effect covalent crosslinking to a target nucleotide using an alkylating agent complementary to the single-stranded target nucleotide sequence. A photoactivated crosslinking to single-stranded oligonucleotides mediated by psoralen was disclosed by Lee, B. L., *et al.*, *Biochemistry* (1988) 27:3197-3203. Use of crosslinking in triple-helix forming probes was also disclosed by Home, *et al.*, *J Am Chem Soc* (1990) 112:2435-2437. Use of N⁴, N⁴-ethanocytosine as an alkylating agent to crosslink to single-stranded oligonucleotides has also been described by Webb and Matteucci, *J Am Chem Soc* (1986) 108:2764-2765; *Nucleic Acids Res* (1986) 14:7661-7674; Feteritz *et al.*, *J. Am. Chem. Soc.* 113:4000 (1991). Various compounds to bind, detect, label, and/or cleave nucleic acids are known in the art. See, for example, U.S. Patent Nos. 5,543,507; 5,672,593; 5,484,908; 5,256,648; and, 5,681,941.

Proteins

The isolated proteins of the present invention comprise a polypeptide having at least 10 amino acids encoded by any one of the polynucleotides of the present invention as discussed more fully, *supra*, or polypeptides which are conservatively modified variants thereof. The proteins of the present invention or variants thereof can comprise any number of contiguous amino acid residues from a polypeptide of the present invention, wherein that number is selected from the group of integers consisting of from 10 to the number of residues in a full-length polypeptide of the present invention. Optionally, this subsequence of contiguous amino acids is at least 15, 20, 25, 30, 35, or 40 amino acids in length, often at least 50, 60, 70, 80, or 90 amino acids in length. Further, the number of such subsequences can be any integer selected from the group consisting of from 1 to 20, such as 2, 3, 4, or 5.

The present invention further provides a protein comprising a polypeptide having a specified sequence identity with a polypeptide of the present invention. The percentage of sequence identity is an integer selected from the group consisting of from 60 to 99. Exemplary sequence identity values include 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, and 99%.

As those of skill will appreciate, the present invention includes catalytically active polypeptides of the present invention (i.e., enzymes). Catalytically active polypeptides have a specific activity of at least 20%, 30%, or 40%, and preferably at least 50%, 60%,

- 46 -

or 70%, and most preferably at least 80%, 90%, or 95% that of the native (non-synthetic), endogenous polypeptide. Further, the substrate specificity (k_{cat}/K_m) is optionally substantially similar to the native (non-synthetic), endogenous polypeptide. Typically, the K_m will be at least 30%, 40%, or 50%, that of the native (non-synthetic), endogenous polypeptide; and more preferably at least 60%, 70%, 80%, or 90%. Methods of assaying and quantifying measures of enzymatic activity and substrate specificity (k_{cat}/K_m), are well known to those of skill in the art.

Generally, the proteins of the present invention will, when presented as an immunogen, elicit production of an antibody specifically reactive to a polypeptide of the present invention. Further, the proteins of the present invention will not bind to antisera raised against a polypeptide of the present invention which has been fully immunosorbed with the same polypeptide. Immunoassays for determining binding are well known to those of skill in the art. A preferred immunoassay is a competitive immunoassay as discussed, *infra*. Thus, the proteins of the present invention can be employed as immunogens for constructing antibodies immunoreactive to a protein of the present invention for such exemplary utilities as immunoassays or protein purification techniques.

Expression of Proteins in Host Cells

Using the nucleic acids of the present invention, one may express a protein of the present invention in a recombinantly engineered cell such as bacteria, yeast, insect, mammalian, or preferably plant cells. The cells produce the protein in a non-natural condition (e.g., in quantity, composition, location, and/or time), because they have been genetically altered through human intervention to do so.

It is expected that those of skill in the art are knowledgeable in the numerous expression systems available for expression of a nucleic acid encoding a protein of the present invention. No attempt to describe in detail the various methods known for the expression of proteins in prokaryotes or eukaryotes will be made.

In brief summary, the expression of isolated nucleic acids encoding a protein of the present invention will typically be achieved by operably linking, for example, the DNA or cDNA to a promoter (which is either constitutive or inducible), followed by incorporation into an expression vector. The vectors can be suitable for replication and integration in either prokaryotes or eukaryotes. Typical expression vectors contain

- 47 -

transcription and translation terminators, initiation sequences, and promoters useful for regulation of the expression of the DNA encoding a protein of the present invention. To obtain high level expression of a cloned gene, it is desirable to construct expression vectors which contain, at the minimum, a strong promoter to direct transcription, a
5 ribosome binding site for translational initiation, and a transcription/translation terminator. One of skill would recognize that modifications can be made to a protein of the present invention without diminishing its biological activity. Some modifications may be made to facilitate the cloning, expression, or incorporation of the targeting molecule into a fusion protein. Such modifications are well known to those of skill in
10 the art and include, for example, a methionine added at the amino terminus to provide an initiation site, or additional amino acids (*e.g.*, poly His) placed on either terminus to create conveniently located restriction sites or termination codons or purification sequences.

15 *A. Expression in Prokaryotes*

Prokaryotic cells may be used as hosts for expression. Prokaryotes most frequently are represented by various strains of *E. coli*; however, other microbial strains may also be used. Commonly used prokaryotic control sequences which are defined herein to include promoters for transcription initiation, optionally with an operator,
20 along with ribosome binding site sequences, include such commonly used promoters as the beta lactamase (penicillinase) and lactose (lac) promoter systems (Chang et al., Nature 198:1056 (1977)), the tryptophan (trp) promoter system (Goeddel et al., Nucleic Acids Res. 8:4057 (1980)) and the lambda derived P_L promoter and N-gene ribosome binding site (Shimatake *et al.*, Nature 292:128 (1981)). The inclusion of selection
25 markers in DNA vectors transfected in *E. coli* is also useful. Examples of such markers include genes specifying resistance to ampicillin, tetracycline, or chloramphenicol.

The vector is selected to allow introduction into the appropriate host cell. Bacterial vectors are typically of plasmid or phage origin. Appropriate bacterial cells are infected with phage vector particles or transfected with naked phage vector DNA. If a
30 plasmid vector is used, the bacterial cells are transfected with the plasmid vector DNA. Expression systems for expressing a protein of the present invention are available using *Bacillus sp.* and *Salmonella* (Palva, *et al.*, Gene 22: 229-235 (1983); Mosbach, *et al.*, Nature 302: 543-545 (1983)).

B. Expression in Eukaryotes

A variety of eukaryotic expression systems such as yeast, insect cell lines, plant and mammalian cells, are known to those of skill in the art. As explained briefly below, a of the present invention can be expressed in these eukaryotic systems. In some
5 embodiments, transformed/transfected plant cells, as discussed *infra*, are employed as expression systems for production of the proteins of the instant invention.

Synthesis of heterologous proteins in yeast is well known. Sherman, F., *et al.*, *Methods in Yeast Genetics*, Cold Spring Harbor Laboratory (1982) is a well recognized work describing the various methods available to produce the protein in yeast. Two
10 widely utilized yeast for production of eukaryotic proteins are *Saccharomyces cerevisiae* and *Pichia pastoris*. Vectors, strains, and protocols for expression in *Saccharomyces* and *Pichia* are known in the art and available from commercial suppliers (e.g., Invitrogen). Suitable vectors usually have expression control sequences, such as promoters, including 3-phosphoglycerate kinase or alcohol oxidase, and an origin of
15 replication, termination sequences and the like as desired.

A protein of the present invention, once expressed, can be isolated from yeast by lysing the cells and applying standard protein isolation techniques to the lysates. The monitoring of the purification process can be accomplished by using Western blot techniques or radioimmunoassay of other standard immunoassay techniques.

20 The sequences encoding proteins of the present invention can also be ligated to various expression vectors for use in transfecting cell cultures of, for instance, mammalian, insect, or plant origin. Illustrative of cell cultures useful for the production of the peptides are mammalian cells. Mammalian cell systems often will be in the form of monolayers of cells although mammalian cell suspensions may also be used. A
25 number of suitable host cell lines capable of expressing intact proteins have been developed in the art, and include the HEK293, BHK21, and CHO cell lines. Expression vectors for these cells can include expression control sequences, such as an origin of replication, a promoter (e.g., the CMV promoter, a HSV *tk* promoter or *pgk* (phosphoglycerate kinase) promoter), an enhancer (Queen *et al.*, *Immunol. Rev.* 89: 49
30 (1986)), and necessary processing information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites (e.g., an SV40 large T Ag poly A addition site), and transcriptional terminator sequences. Other animal cells useful for production of

proteins of the present invention are available, for instance, from the American Type Culture Collection Catalogue of Cell Lines and Hybridomas (7th edition, 1992).

Appropriate vectors for expressing proteins of the present invention in insect cells are usually derived from the SF9 baculovirus. Suitable insect cell lines include mosquito
5 larvae, silkworm, armyworm, moth and *Drosophila* cell lines such as a Schneider cell line (See Schneider, *J. Embryol. Exp. Morphol.* 27: 353-365 (1987).

As with yeast, when higher animal or plant host cells are employed, polyadenylation or transcription terminator sequences are typically incorporated into the vector. An example of a terminator sequence is the polyadenylation sequence from the
10 bovine growth hormone gene. Sequences for accurate splicing of the transcript may also be included. An example of a splicing sequence is the VP1 intron from SV40 (Sprague, *et al.*, *J. Virol.* 45: 773-781 (1983)). Additionally, gene sequences to control replication in the host cell may be incorporated into the vector such as those found in bovine papilloma virus type-vectors. Saveria-Campo, M., Bovine Papilloma Virus DNA a
15 Eukaryotic Cloning Vector in *DNA Cloning Vol. II a Practical Approach*, D.M. Glover, Ed., IRL Press, Arlington, Virginia pp. 213-238 (1985).

Transfection/Transformation of Cells

The method of transformation/transfection is not critical to the instant invention;
20 various methods of transformation or transfection are currently available. As newer methods are available to transform crops or other host cells they may be directly applied. Accordingly, a wide variety of methods have been developed to insert a DNA sequence into the genome of a host cell to obtain the transcription and/or translation of the sequence to effect phenotypic changes in the organism. Thus, any method which
25 provides for efficient transformation/transfection may be employed.

A. Plant Transformation

A DNA sequence coding for the desired polynucleotide of the present invention, for example a cDNA or a genomic sequence encoding a full length protein, will be used
30 to construct a recombinant expression cassette which can be introduced into the desired plant.

Isolated nucleic acid acids of the present invention can be introduced into plants according techniques known in the art. Generally, recombinant expression cassettes as

described above and suitable for transformation of plant cells are prepared. Techniques for transforming a wide variety of higher plant species are well known and described in the technical, scientific, and patent literature. See, for example, Weising *et al.*, *Ann. Rev. Genet.* 22: 421-477 (1988). For example, the DNA construct may be introduced
5 directly into the genomic DNA of the plant cell using techniques such as electroporation, PEG poration, particle bombardment, silicon fiber delivery, or microinjection of plant cell protoplasts or embryogenic callus. See, e.g., Tomes, *et al.*, Direct DNA Transfer into Intact Plant Cells Via Microprojectile Bombardment. pp.197-213 in Plant Cell, Tissue and Organ Culture, Fundamental Methods. eds. O. L. Gamborg and G.C.
10 Phillips. Springer-Verlag Berlin Heidelberg New York, 1995. Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is infected by the bacteria. See, U.S.
15 Patent No. 5,591,616.

The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski *et al.*, *Embo J.* 3: 2717-2722 (1984). Electroporation techniques are described in Fromm *et al.*, *Proc. Natl. Acad. Sci.* 82: 5824 (1985). Ballistic transformation techniques are described in Klein *et al.*, *Nature* 327: 70-73
20 (1987). *Agrobacterium tumefaciens*-mediated transformation techniques are well described in the scientific literature. See, for example Horsch *et al.*, *Science* 233: 496-498 (1984), and Fraley *et al.*, *Proc. Natl. Acad. Sci.* 80: 4803 (1983). Although *Agrobacterium* is useful primarily in dicots, certain monocots can be transformed by *Agrobacterium*. For instance, *Agrobacterium* transformation of maize is described in
25 U.S. Patent No. 5,550,318.

Other methods of transfection or transformation include (1) *Agrobacterium rhizogenes*-mediated transformation (see, e.g., Lichtenstein and Fuller In: Genetic Engineering, vol. 6, PWJ Rigby, Ed., London, Academic Press, 1987; and Lichtenstein, C. P., and Draper, J., In: DNA Cloning, Vol. II, D. M. Glover, Ed., Oxford, IRI
30 Press, 1985), Application PCT/US87/02512 (WO 88/02405 published Apr. 7, 1988) describes the use of *A. rhizogenes* strain A4 and its Ri plasmid along with *A. tumefaciens* vectors pARC8 or pARC16 (2) liposome-mediated DNA uptake (see, e.g., Freeman *et*

al., Plant Cell Physiol. 25: 1353, 1984), (3) the vortexing method (see, e.g., Kindle, *Proc. Natl. Acad. Sci.*, USA 87: 1228, (1990).

DNA can also be introduced into plants by direct DNA transfer into pollen as described by Zhou *et al.*, Methods in Enzymology, 101:433 (1983); D. Hess, Intern
5 Rev. Cytol., 107:367 (1987); Luo *et al.*, Plant Mol. Biol. Reporter, 6:165 (1988).
Expression of polypeptide coding genes can be obtained by injection of the DNA into
reproductive organs of a plant as described by Pena *et al.*, Nature, 325:274 (1987).
DNA can also be injected directly into the cells of immature embryos and the
rehydration of desiccated embryos as described by Neuhaus *et al.*, Theor. Appl. Genet.,
10 75:30 (1987); and Benbrook *et al.*, in Proceedings Bio Expo 1986, Butterworth,
Stoneham, Mass., pp. 27-54 (1986). A variety of plant viruses that can be employed as
vectors are known in the art and include cauliflower mosaic virus (CaMV), geminivirus,
brome mosaic virus, and tobacco mosaic virus.

15 *B. Transfection of Prokaryotes, Lower Eukaryotes, and Animal Cells*

Animal and lower eukaryotic (e.g., yeast) host cells are competent or rendered
competent for transfection by various means. There are several well-known methods of
introducing DNA into animal cells. These include: calcium phosphate precipitation,
fusion of the recipient cells with bacterial protoplasts containing the DNA, treatment of
20 the recipient cells with liposomes containing the DNA, DEAE dextran, electroporation,
biolistics, and micro-injection of the DNA directly into the cells. The transfected cells
are cultured by means well known in the art. Kuchler, R.J., *Biochemical Methods in Cell
Culture and Virology*, Dowden, Hutchinson and Ross, Inc. (1977).

25 Synthesis of Proteins

The proteins of the present invention can be constructed using non-cellular
synthetic methods. Solid phase synthesis of proteins of less than about 50 amino acids in
length may be accomplished by attaching the C-terminal amino acid of the sequence to an
insoluble support followed by sequential addition of the remaining amino acids in the
30 sequence. Techniques for solid phase synthesis are described by Barany and Merrifield,
Solid-Phase Peptide Synthesis, pp. 3-284 in *The Peptides: Analysis, Synthesis, Biology*.
Vol. 2: Special Methods in Peptide Synthesis, Part A.; Merrifield, *et al.*, *J. Am. Chem.*
Soc. 85: 2149-2156 (1963), and Stewart *et al.*, *Solid Phase Peptide Synthesis*, 2nd ed.,

Pierce Chem. Co., Rockford, Ill. (1984). Proteins of greater length may be synthesized by condensation of the amino and carboxy termini of shorter fragments. Methods of forming peptide bonds by activation of a carboxy terminal end (e.g., by the use of the coupling reagent N,N'-dicyclohexylcarbodiimide)) is known to those of skill.

5

Purification of Proteins

The proteins of the present invention may be purified by standard techniques well known to those of skill in the art. Recombinantly produced proteins of the present invention can be directly expressed or expressed as a fusion protein. The recombinant protein is purified by a combination of cell lysis (e.g., sonication, French press) and affinity chromatography. For fusion products, subsequent digestion of the fusion protein with an appropriate proteolytic enzyme releases the desired recombinant protein.

The proteins of this invention, recombinant or synthetic, may be purified to substantial purity by standard techniques well known in the art, including detergent solubilization, selective precipitation with such substances as ammonium sulfate, column chromatography, immunopurification methods, and others. See, for instance, R. Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag: New York (1982); Deutscher, *Guide to Protein Purification*, Academic Press (1990). For example, antibodies may be raised to the proteins as described herein. Purification from *E. coli* can be achieved following procedures described in U.S. Patent No. 4,511,503. The protein may then be isolated from cells expressing the protein and further purified by standard protein chemistry techniques as described herein. Detection of the expressed protein is achieved by methods known in the art and include, for example, radioimmunoassays, Western blotting techniques or immunoprecipitation.

25

Transgenic Plant Regeneration

Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed genotype. Such regeneration techniques often rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker which has been introduced together with a polynucleotide of the present invention. For transformation and regeneration of maize see, Gordon-Kamm *et al.*, *The Plant Cell*, 2:603-618 (1990).

30

Plants cells transformed with a plant expression vector can be regenerated, e.g., from single cells, callus tissue or leaf discs according to standard plant tissue culture techniques. It is well known in the art that various cells, tissues, and organs from almost any plant can be successfully cultured to regenerate an entire plant. Plant regeneration
5 from cultured protoplasts is described in Evans *et al.*, *Protoplasts Isolation and Culture, Handbook of Plant Cell Culture*, Macmillilan Publishing Company, New York, pp. 124-176 (1983); and Binding, *Regeneration of Plants, Plant Protoplasts*, CRC Press, Boca Raton, pp. 21-73 (1985).

The regeneration of plants containing the foreign gene introduced by
10 *Agrobacterium* from leaf explants can be achieved as described by Horsch *et al.*, *Science*, 227:1229-1231 (1985). In this procedure, transformants are grown in the presence of a selection agent and in a medium that induces the regeneration of shoots in the plant species being transformed as described by Fraley *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 80:4803 (1983). This procedure typically produces shoots within two to four
15 weeks and these transformant shoots are then transferred to an appropriate root-inducing medium containing the selective agent and an antibiotic to prevent bacterial growth. Transgenic plants of the present invention may be fertile or sterile.

Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee *et al.*, *Ann. Rev. of Plant Phys.* 38: 467-486 (1987). The regeneration of plants from either single plant
20 protoplasts or various explants is well known in the art. See, for example, *Methods for Plant Molecular Biology*, A. Weissbach and H. Weissbach, eds., Academic Press, Inc., San Diego, Calif. (1988). This regeneration and growth process includes the steps of selection of transformant cells and shoots, rooting the transformant shoots and growth of
25 the plantlets in soil. For maize cell culture and regeneration see generally, *The Maize Handbook*, Freeling and Walbot, Eds., Springer, New York (1994); *Corn and Corn Improvement*, 3rd edition, Sprague and Dudley Eds., American Society of Agronomy, Madison, Wisconsin (1988).

One of skill will recognize that after the recombinant expression cassette is stably
30 incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

In vegetatively propagated crops, mature transgenic plants can be propagated by the taking of cuttings or by tissue culture techniques to produce multiple identical plants. Selection of desirable transgenics is made and new varieties are obtained and propagated vegetatively for commercial use. In seed propagated crops, mature transgenic plants can
5 be self crossed to produce a homozygous inbred plant. The inbred plant produces seed containing the newly introduced heterologous nucleic acid. These seeds can be grown to produce plants that would produce the selected phenotype.

Parts obtained from the regenerated plant, such as flowers, seeds, leaves, branches, fruit, and the like are included in the invention, provided that these parts
10 comprise cells comprising the isolated nucleic acid of the present invention. Progeny and variants, and mutants of the regenerated plants are also included within the scope of the invention, provided that these parts comprise the introduced nucleic acid sequences.

Transgenic plants expressing the selectable marker can be screened for transmission of the nucleic acid of the present invention by, for example, standard
15 immunoblot and DNA detection techniques. Transgenic lines are also typically evaluated on levels of expression of the heterologous nucleic acid. Expression at the RNA level can be determined initially to identify and quantitate expression-positive plants. Standard techniques for RNA analysis can be employed and include PCR amplification assays using oligonucleotide primers designed to amplify only the heterologous RNA templates
20 and solution hybridization assays using heterologous nucleic acid-specific probes. The RNA-positive plants can then analyzed for protein expression by Western immunoblot analysis using the specifically reactive antibodies of the present invention. In addition, *in situ* hybridization and immunocytochemistry according to standard protocols can be done using heterologous nucleic acid specific polynucleotide probes and antibodies,
25 respectively, to localize sites of expression within transgenic tissue. Generally, a number of transgenic lines are usually screened for the incorporated nucleic acid to identify and select plants with the most appropriate expression profiles.

A preferred embodiment is a transgenic plant that is homozygous for the added heterologous nucleic acid; i.e., a transgenic plant that contains two added nucleic acid
30 sequences, one gene at the same locus on each chromosome of a chromosome pair. A homozygous transgenic plant can be obtained by sexually mating (selfing) a heterozygous transgenic plant that contains a single added heterologous nucleic acid, germinating some of the seed produced and analyzing the resulting plants produced for altered expression

of a polynucleotide of the present invention relative to a control plant (i.e., native, non-transgenic). Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated.

5 Modulating Polypeptide Levels and/or Composition

The present invention further provides a method for modulating (i.e., increasing or decreasing) the concentration or composition of the polypeptides of the present invention in a plant or part thereof. Modulation can be effected by increasing or decreasing the concentration and/or the composition (i.e., the ratio of the polypeptides of
10 the present invention) in a plant. The method comprises transforming a plant cell with a recombinant expression cassette comprising a polynucleotide of the present invention as described above to obtain a transformed plant cell, growing the transformed plant cell under plant forming conditions, and inducing expression of a polynucleotide of the present invention in the plant for a time sufficient to modulate concentration and/or
15 composition in the plant or plant part.

In some embodiments, the content and/or composition of polypeptides of the present invention in a plant may be modulated by altering, *in vivo* or *in vitro*, the promoter of a non-isolated gene of the present invention to up- or down-regulate gene expression. In some embodiments, the coding regions of native genes of the present
20 invention can be altered via substitution, addition, insertion, or deletion to decrease activity of the encoded enzyme. See, e.g., Kmiec, U.S. Patent 5,565,350; Zarling *et al.*, PCT/US93/03868. And in some embodiments, an isolated nucleic acid (e.g., a vector) comprising a promoter sequence is transfected into a plant cell. Subsequently, a plant cell comprising the promoter operably linked to a polynucleotide of the present invention
25 is selected for by means known to those of skill in the art such as, but not limited to, Southern blot, DNA sequencing, or PCR analysis using primers specific to the promoter and to the gene and detecting amplicons produced therefrom. A plant or plant part altered or modified by the foregoing embodiments is grown under plant forming conditions for a time sufficient to modulate the concentration and/or composition of
30 polypeptides of the present invention in the plant. Plant forming conditions are well known in the art and discussed briefly, *supra*.

In general, concentration or composition is increased or decreased by at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% relative to a native control

- 56 -

plant, plant part, or cell lacking the aforementioned recombinant expression cassette. Modulation in the present invention may occur during and/or subsequent to growth of the plant to the desired stage of development. Modulating nucleic acid expression temporally and/or in particular tissues can be controlled by employing the appropriate promoter operably linked to a polynucleotide of the present invention in, for example, sense or antisense orientation as discussed in greater detail, *supra*. Induction of expression of a polynucleotide of the present invention can also be controlled by exogenous administration of an effective amount of inducing compound. Inducible promoters and inducing compounds which activate expression from these promoters are well known in the art. In preferred embodiments, the polypeptides of the present invention are modulated in monocots, particularly maize.

Molecular Markers

The present invention provides a method of genotyping a plant comprising a polynucleotide of the present invention. Preferably, the plant is a monocot, such as maize or sorghum. Genotyping provides a means of distinguishing homologs of a chromosome pair and can be used to differentiate segregants in a plant population. Molecular marker methods can be used for phylogenetic studies, characterizing genetic relationships among crop varieties, identifying crosses or somatic hybrids, localizing chromosomal segments affecting monogenic traits, map based cloning, and the study of quantitative inheritance. See, e.g., *Plant Molecular Biology: A Laboratory Manual*, Chapter 7, Clark, Ed., Springer-Verlag, Berlin (1997). For molecular marker methods, see generally, *The DNA Revolution* by Andrew H. Paterson 1996 (Chapter 2) in: *Genome Mapping in Plants* (ed. Andrew H. Paterson) by Academic Press/R. G. Landis Company, Austin, Texas, pp.7-21.

The particular method of genotyping in the present invention may employ any number of molecular marker analytic techniques such as, but not limited to, restriction fragment length polymorphisms (RFLPs). RFLPs are the product of allelic differences between DNA restriction fragments caused by nucleotide sequence variability. As is well known to those of skill in the art, RFLPs are typically detected by extraction of genomic DNA and digestion with a restriction enzyme. Generally, the resulting fragments are separated according to size and hybridized with a probe; single copy probes are preferred. Restriction fragments from homologous chromosomes are

revealed. Differences in fragment size among alleles represent an RFLP. Thus, the present invention further provides a means to follow segregation of a gene or nucleic acid of the present invention as well as chromosomal sequences genetically linked to these genes or nucleic acids using such techniques as RFLP analysis. Linked
5 chromosomal sequences are within 50 centiMorgans (cM), often within 40 or 30 cM, preferably within 20 or 10 cM, more preferably within 5, 3, 2, or 1 cM of a gene of the present invention.

In the present invention, the nucleic acid probes employed for molecular marker mapping of plant nuclear genomes selectively hybridize, under selective hybridization
10 conditions, to a gene encoding a polynucleotide of the present invention. In preferred embodiments, the probes are selected from polynucleotides of the present invention. Typically, these probes are cDNA probes or *Pst I* genomic clones. The length of the probes is discussed in greater detail, *supra*, but are typically at least 15 bases in length, more preferably at least 20, 25, 30, 35, 40, or 50 bases in length. Generally, however,
15 the probes are less than about 1 kilobase in length. Preferably, the probes are single copy probes that hybridize to a unique locus in a haploid chromosome complement. Some exemplary restriction enzymes employed in RFLP mapping are EcoRI, EcoRV, and SstI. As used herein the term "restriction enzyme" includes reference to a composition that recognizes and, alone or in conjunction with another composition,
20 cleaves at a specific nucleotide sequence.

The method of detecting an RFLP comprises the steps of (a) digesting genomic DNA of a plant with a restriction enzyme; (b) hybridizing a nucleic acid probe, under selective hybridization conditions, to a sequence of a polynucleotide of the present of said genomic DNA; (c) detecting therefrom a RFLP. Other methods of differentiating
25 polymorphic (allelic) variants of polynucleotides of the present invention can be had by utilizing molecular marker techniques well known to those of skill in the art including such techniques as: 1) single stranded conformation analysis (SSCP); 2) denaturing gradient gel electrophoresis (DGGE); 3) RNase protection assays; 4) allele-specific oligonucleotides (ASOs); 5) the use of proteins which recognize nucleotide mismatches, such as the *E. coli* mutS protein; and 6) allele-specific PCR. Other approaches based on
30 the detection of mismatches between the two complementary DNA strands include clamped denaturing gel electrophoresis (CDGE); heteroduplex analysis (HA); and chemical mismatch cleavage (CMC). Exemplary polymorphic variants are provided in

Table I, *supra*. Thus, the present invention further provides a method of genotyping comprising the steps of contacting, under stringent hybridization conditions, a sample suspected of comprising a polynucleotide of the present invention with a nucleic acid probe. Generally, the sample is a plant sample; preferably, a sample suspected of comprising a maize polynucleotide of the present invention (e.g., gene, mRNA). The nucleic acid probe selectively hybridizes, under stringent conditions, to a subsequence of a polynucleotide of the present invention comprising a polymorphic marker. Selective hybridization of the nucleic acid probe to the polymorphic marker nucleic acid sequence yields a hybridization complex. Detection of the hybridization complex indicates the presence of that polymorphic marker in the sample. In preferred embodiments, the nucleic acid probe comprises a polynucleotide of the present invention.

UTR's and Codon Preference

In general, translational efficiency has been found to be regulated by specific sequence elements in the 5' non-coding or untranslated region (5' UTR) of the RNA. Positive sequence motifs include translational initiation consensus sequences (Kozak, *Nucleic Acids Res.* 15:8125 (1987)) and the 7-methylguanosine cap structure (Drummond *et al.*, *Nucleic Acids Res.* 13:7375 (1985)). Negative elements include stable intramolecular 5' UTR stem-loop structures (Muesing *et al.*, *Cell* 48:691 (1987)) and AUG sequences or short open reading frames preceded by an appropriate AUG in the 5' UTR (Kozak, *supra*, Rao *et al.*, *Mol. and Cell. Biol.* 8:284 (1988)). Accordingly, the present invention provides 5' and/or 3' UTR regions for modulation of translation of heterologous coding sequences.

Further, the polypeptide-encoding segments of the polynucleotides of the present invention can be modified to alter codon usage. Altered codon usage can be employed to alter translational efficiency and/or to optimize the coding sequence for expression in a desired host or to optimize the codon usage in a heterologous sequence for expression in maize. Codon usage in the coding regions of the polynucleotides of the present invention can be analyzed statistically using commercially available software packages such as "Codon Preference" available from the University of Wisconsin Genetics Computer Group (see Devereaux *et al.*, *Nucleic Acids Res.* 12: 387-395 (1984)) or MacVector 4.1 (Eastman Kodak Co., New Haven, Conn.). Thus, the present invention provides a codon usage frequency characteristic of the coding region of at least one of

the polynucleotides of the present invention. The number of polynucleotides that can be used to determine a codon usage frequency can be any integer from 1 to the number of polynucleotides of the present invention as provided herein. Optionally, the polynucleotides will be full-length sequences. An exemplary number of sequences for statistical analysis can be at least 1, 5, 10, 20, 50, or 100.

Sequence Shuffling

The present invention provides methods for sequence shuffling using polynucleotides of the present invention, and compositions resulting therefrom.

Sequence shuffling is described in PCT publication No. WO 97/20078. See also, Zhang, J.- H., *et al. Proc. Natl. Acad. Sci. USA* 94:4504-4509 (1997). Generally, sequence shuffling provides a means for generating libraries of polynucleotides having a desired characteristic which can be selected or screened for. Libraries of recombinant polynucleotides are generated from a population of related sequence polynucleotides which comprise sequence regions which have substantial sequence identity and can be homologously recombined in vitro or in vivo. The population of sequence-recombined polynucleotides comprises a subpopulation of polynucleotides which possess desired or advantageous characteristics and which can be selected by a suitable selection or screening method. The characteristics can be any property or attribute capable of being selected for or detected in a screening system, and may include properties of: an encoded protein, a transcriptional element, a sequence controlling transcription, RNA processing, RNA stability, chromatin conformation, translation, or other expression property of a gene or transgene, a replicative element, a protein-binding element, or the like, such as any feature which confers a selectable or detectable property. In some embodiments, the selected characteristic will be a decreased K_m and/or increased K_{cat} over the wild-type protein as provided herein. In other embodiments, a protein or polynucleotide generated from sequence shuffling will have a ligand binding affinity greater than the non-shuffled wild-type polynucleotide. The increase in such properties can be at least 110%, 120%, 130%, 140% or at least 150% of the wild-type value.

Assays for Compounds that Modulate Enzymatic Activity or Expression

The present invention also provides means for identifying compounds that bind to (e.g., substrates), and/or increase or decrease (i.e., modulate) the enzymatic activity of, catalytically active polypeptides of the present invention. The method comprises

5 contacting a polypeptide of the present invention with a compound whose ability to bind to or modulate enzyme activity is to be determined. The polypeptide employed will have at least 20%, preferably at least 30% or 40%, more preferably at least 50% or 60%, and most preferably at least 70% or 80% of the specific activity of the native, full-length polypeptide of the present invention (e.g., enzyme). Generally, the polypeptide will be

10 present in a range sufficient to determine the effect of the compound, typically about 1 nM to 10 μ M. Likewise, the compound will be present in a concentration of from about 1 nM to 10 μ M. Those of skill will understand that such factors as enzyme concentration, ligand concentrations (i.e., substrates, products, inhibitors, activators), pH, ionic strength, and temperature will be controlled so as to obtain useful kinetic data

15 and determine the presence or absence of a compound that binds or modulates polypeptide activity. Methods of measuring enzyme kinetics is well known in the art. See, e.g., Segel, *Biochemical Calculations*, 2nd ed., John Wiley and Sons, New York (1976).

Although the present invention has been described in some detail by way of

20 illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

Example 1

25 This example describes the construction of the cDNA libraries.

Total RNA Isolation

Total RNA was isolated from corn tissues with TRIzol Reagent (Life Technology Inc. Gaithersburg, MD) using a modification of the guanidine isothiocyanate/acid-phenol

30 procedure described by Chomczynski and Sacchi (Chomczynski, P., and Sacchi, N. *Anal. Biochem.* 162, 156 (1987)). In brief, plant tissue samples were pulverized in liquid nitrogen before the addition of the TRIzol Reagent, and then were further homogenized with a mortar and pestle. Addition of chloroform followed by

- 61 -

centrifugation was conducted for separation of an aqueous phase and an organic phase. The total RNA was recovered by precipitation with isopropyl alcohol from the aqueous phase.

5 Poly(A)+ RNA Isolation

The selection of poly(A)+ RNA from total RNA was performed using PolyATtract system (Promega Corporation, Madison, WI). In brief, biotinylated oligo(dT) primers were used to hybridize to the 3' poly(A) tails on mRNA. The hybrids were captured using streptavidin coupled to paramagnetic particles and a magnetic
10 separation stand. The mRNA was washed at high stringent condition and eluted by RNase-free deionized water.

cDNA Library Construction

cDNA synthesis was performed and unidirectional cDNA libraries were
15 constructed using the SuperScript Plasmid System (Life Technology Inc. Gaithersburg, MD). The first strand of cDNA was synthesized by priming an oligo(dT) primer containing a Not I site. The reaction was catalyzed by SuperScript Reverse Transcriptase II at 45°C. The second strand of cDNA was labeled with alpha-³²P-dCTP and a portion of the reaction was analyzed by agarose gel electrophoresis to determine
20 cDNA sizes. cDNA molecules smaller than 500 base pairs and unligated adapters were removed by Sephacryl-S400 chromatography. The selected cDNA molecules were ligated into pSPORT1 vector in between of Not I and Sal I sites.

Example 2

25 This example describes cDNA sequencing and library subtraction.

Sequencing Template Preparation

Individual colonies were picked and DNA was prepared either by PCR with M13 forward primers and M13 reverse primers, or by plasmid isolation. All the cDNA
30 clones were sequenced using M13 reverse primers.

Q-bot Subtraction Procedure

cDNA libraries subjected to the subtraction procedure were plated out on 22 x 22 cm² agar plate at density of about 3,000 colonies per plate. The plates were incubated in a 37°C incubator for 12-24 hours. Colonies were picked into 384-well plates by a robot colony picker, Q-bot (GENETIX Limited). These plates were incubated overnight at 37°C.

Once sufficient colonies were picked, they were pinned onto 22 x 22 cm² nylon membranes using Q-bot. Each membrane contained 9,216 colonies or 36,864 colonies. These membranes were placed onto agar plate with appropriate antibiotic. The plates were incubated at 37°C for overnight.

After colonies were recovered on the second day, these filters were placed on filter paper prewetted with denaturing solution for four minutes, then were incubated on top of a boiling water bath for additional four minutes. The filters were then placed on filter paper prewetted with neutralizing solution for four minutes. After excess solution was removed by placing the filters on dry filter papers for one minute, the colony side of the filters were placed into Proteinase K solution, incubated at 37°C for 40-50 minutes. The filters were placed on dry filter papers to dry overnight. DNA was then cross-linked to nylon membrane by UV light treatment.

Colony hybridization was conducted as described by Sambrook, J., Fritsch, E.F. and Maniatis, T., (in *Molecular Cloning: A laboratory Manual*, 2nd Edition). The following probes were used in colony hybridization:

1. First strand cDNA from the same tissue as the library was made from to remove the most redundant clones.
2. 48-192 most redundant cDNA clones from the same library based on previous sequencing data.
3. 192 most redundant cDNA clones in the corn sequence database.
4. A Sal-A20 oligo nucleotide: TCG ACC CAC GCG TCC GAA AAA AAA AAA AAA AAA AAA, listed in SEQ ID NO: 3, removes clones containing a poly A tail but no cDNA.
5. cDNA clones derived from rRNA.

The image of the autoradiography was scanned into computer and the signal intensity and cold colony addresses of each colony was analyzed. Re-arraying of cold-colonies from 384 well plates to 96 well plates was conducted using Q-bot.

5 Example 3

This example describes identification of the gene from a computer homology search.

Gene identities were determined by conducting BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al., (1990) *J. Mol. Biol.* 215:403-410; see also
10 www.ncbi.nlm.nih.gov/BLAST/) searches under default parameters for similarity to sequences contained in the BLAST "nr" database (comprising all non-redundant GenBank CDS translations, sequences derived from the 3-dimensional structure Brookhaven Protein Data Bank, the last major release of the SWISS-PROT protein sequence database, EMBL, and DDBJ databases). The cDNA sequences were analyzed
15 for similarity to all publicly available DNA sequences contained in the "nr" database using the BLASTN algorithm. The DNA sequences were translated in all reading frames and compared for similarity to all publicly available protein sequences contained in the "nr" database using the BLASTX algorithm (Gish, W. and States, D. J. (1993) *Nature Genetics* 3:266-272) provided by the NCBI. In some cases, the sequencing data
20 from two or more clones containing overlapping segments of DNA were used to construct contiguous DNA sequences.

SEQ ID NO:1 is the full length cDNA sequence of the maize orthologue of Ku70 obtained from a library constructed using root tissue from corn root worm infested B73 at the V5 stage. The sequence was generated from a partial sequence encoding
25 approximately 70% of the C-terminal end. Reverse-transcriptase PCR generated the remaining N-terminal end. This 2.4kb sequence contains an open reading frame encoding a protein of 73kDa. The deduced amino acid sequence of this protein indicates that there are 51 aspartic acid and 46 glutamic acid residues, a characteristic of the acidic nature of all Ku homologues studied to date. A comparison of the maize Ku70 deduced amino acid
30 sequence with that from human (Accession No. J04611), mouse (Accession No. P23475), chicken (Accession No. O93257), and *D. melanogaster* (Accession No. Q26228) proteins shows about a 40-42% similarity between maize Ku70 and vertebrate Ku70.

References

1. Friedburg, E., Walker, G. and Siede, W. (1995) DNA Repair and Mutagenesis
ASM Press, Washington DC.
- 5 2. Nickoloff J. and Hoekstra, M. (1998) DNA Damage and Repair.
Humana Press, Totowa, NJ
3. Shinohara and Ogawa (1995) Trends in Biochem. Sci. 237, 387-391
- 10 4. Hays et al. (1995) Proc. Natl. Acad. Sci. USA 92, 6925-6929
5. Lindahl et al. (1995) Trends in Biochem. Sci. 237, 405-411
- 15 6. Jackson & Jeggo, 1995 Trends in Biochem. Sci. 237, 412-415
7. Shah et al. (1995) Anal. Biochem. 227, 1-13
8. Dvir et al. (1992) Proc. Natl. Acad. Sci. USA 89, 11920-11924
- 20 9. Anderson et al. Crit. Rev. Eukaryot. Gene Express. (1992) 4, 283-314
10. Jeggo (1997) Mutation Res. 384, 1-14
- 25 11. Gotlib and Jackson (1993) Cell 72, 131-142
12. Mimori et al. (1981) J. Clin. Invest. 68, 611-620
13. Mimori et al. (1986) J. Biol. Chem. 261, 2274-2278
- 30 14. Yaneva & Arnett (1989) Clin. Exp. Immunol. 76, 366-372
15. Chu (1997) J. Biol. Chem. 272, 24097-24100
- 35 16. Yoo & Dynan (1998) Biochemistry 37, 1336-1343
17. Boulton & Jackson (1996) EMBO J. 15, 5093-5103
18. Roth et al. (1995) Current Biol. 5, 496-498
- 40 19. Casellas et al. (1998) EMBO J. 17, 2404-2411
20. Boulton & Jackson, (1998) EMBO J. 17, 1819-1828
- 45 21. Polotnianska et al. (1998) Current Biol. 8, 831-834
22. Ramsden & Gellert, (1998) EMBO J. 17, 609-614
23. Yang et al., (1996) Mol. Cell. Biol. 16, 3799-3806

24. Burgman et al. (1997) *Cancer Res.* 57, 2847-2850
25. Reeves & Sthoeger (1989) *J. Biol. Chem.* 264, 5047-5052
- 5 26. Chan et al. (1989) *J. Biol. Chem.* 264, 3651-3654
27. Porges et al. (1990) *J. Immunol.* 145, 4222-4228
- 10 28. Jacoby et al. (1994) *J. Biol. Chem.* 269, 11484-11491
29. Paesen et al. (1996) *Biochim Biophys. Acta* 1305, 120-124
30. Boulton & Jackson (1996) *Nucleic Acid Res.* 24, 4639-4648
- 15 31. Feldmann & Winnacker (1993) *J. Biol. Chem.* 268, 12895-12900
32. Feldmann et al. (1996) *J. Biol. Chem.* 271, 27765-27769
- 20 33. Boulton & Jackson (1996) *Nucleic Acid Res.* 24, 4639-4648
34. Wang et al. (1998) *J. Biol. Chem.* 273, 842-848

The above examples are provided to illustrate the invention but not to limit its
25 scope. Other variants of the invention will be readily apparent to one of ordinary skill in
the art and are encompassed by the appended claims. All publications, patents, and
patent applications cited herein are hereby incorporated by reference.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid comprising a member selected from the group consisting of:
 - a) a polynucleotide having at least 60% sequence identity to a polynucleotide encoding the polypeptide of SEQ ID NO: 2;
 - 5 b) a polynucleotide encoding the polypeptide of SEQ ID NO: 2;
 - c) a polynucleotide amplified from a *Zea mays* nucleic acid library using primers which selectively hybridize, under stringent hybridization conditions, to loci within the polynucleotide of SEQ ID NO: 1;
 - d) a polynucleotide which selectively hybridizes, under stringent hybridization
10 conditions and a wash in 2X SSC at 50°C, to the polynucleotide of SEQ ID NO: 1;
 - e) the polynucleotide of SEQ ID NO: 1;
 - f) a polynucleotide which is complementary to a polynucleotide of (a), (b), (c), (d), or (e); and
 - 15 g) a polynucleotide comprising at least 25 contiguous nucleotides from a polynucleotide of (a), (b), (c), (d), (e), or (f).
- 2) A recombinant expression cassette, comprising a member of claim 1 operably linked, in sense or anti-sense orientation, to a promoter.
20
- 3) A host cell comprising the recombinant expression cassette of claim 2.
- 4) A transgenic plant comprising a recombinant expression cassette of claim 2.
- 25 5) The transgenic plant of claim 4, wherein said plant is a monocot.
- 6) The transgenic plant of claim 4, wherein said plant is a dicot.
- 7) The transgenic plant of claim 4, wherein said plant is selected from the group consisting of: maize, soybean, sunflower, sorghum, canola, wheat, alfalfa, cotton,
30 rice, barley, and millet.
- 8) A transgenic seed from the transgenic plant of claim 4.

- 67 -

- 9) A method of modulating the level of Ku70 in a plant, comprising:
- (a) introducing into a plant cell a recombinant expression cassette comprising a Ku70 polynucleotide of claim 1 operably linked to a promoter;
 - (b) culturing the plant cell under plant cell growing conditions; and
 - 5 (c) inducing expression of said polynucleotide for a time sufficient to modulate the level of Ku70 in said plant.
- 10) The method of claim 9, wherein the plant is maize.
- 10 11) An isolated protein comprising a member selected from the group consisting of:
- (a) a polypeptide of at least 20 contiguous amino acids from the polypeptide of SEQ ID NO: 2;
 - (b) the polypeptide of SEQ ID NO: 2;
 - (c) a polypeptide comprising an amino acid sequence which has at least 60%
 - 15 sequence identity to the polypeptide of SEQ ID NO:2;
 - (d) at least one polypeptide encoded by a member of claim 1.

- 1 -

SEQUENCE LISTING

<110> Pioneer Hi-Bred Int'l Inc.

<120> Maize Ku70 Orthologue and Uses Thereof

<130> 0932-PCT

<150> 60/098,986

<151> 1998-09-02

<160> 3

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 2399

<212> DNA

<213> Zea mays

<220>

<221> CDS

<222> (232) ... (2181)

<400> 1

```

gggcagcagc gccgtagagc cgtggtttcc cacaaaaaaaa acggcgccac ggtagcgagt      60
ttctagaacc ttcccacgcc ctgcgcccct ttgctgtctc atacctgtct taaaccctag      120
ctagtttcta ggaccttccc acgccctgcg cccctttgct gtctcaaacc taccttaaac      180
ccttgttagt ttctagaacc ttccagacct ccagttcccg acctcggcgc c atg gac      237
                                         Met Asp
                                         1

ctg gac cca gag ggg atc ttc cgc gac gac agc gat gaa gac gaa gac      285
Leu Asp Pro Glu Gly Ile Phe Arg Asp Asp Ser Asp Glu Asp Glu Asp
      5              10              15

agt gtc cag gag agg gag gcg aac aag gag atg gtt gtc tac ctt gtc      333
Ser Val Gln Glu Arg Glu Ala Asn Lys Glu Met Val Val Tyr Leu Val
      20              25              30

gac gcc tcg ccc aaa atg ttc acc cct gcc acc acc cag gac aat gaa      381
Asp Ala Ser Pro Lys Met Phe Thr Pro Ala Thr Thr Gln Asp Asn Glu
      35              40              45              50

aag cag gag aca cat ttc cat acc att gtt aac tgc atc aca gag tct      429
Lys Gln Glu Thr His Phe His Thr Ile Val Asn Cys Ile Thr Glu Ser
              55              60              65

ctg aag aca caa att att ggt aga tct tat gat gaa gtc gca ata tgt      477
Leu Lys Thr Gln Ile Ile Gly Arg Ser Tyr Asp Glu Val Ala Ile Cys
      70              75              80

ttc ttt aac act aaa gag aag aag aat tta cag gac tca gcc ggt gtt      525
Phe Phe Asn Thr Lys Glu Lys Lys Asn Leu Gln Asp Ser Ala Gly Val
      85              90              95

```

-2-

tat gtt tat aat gtc gga gac aga gag caa ctt gat aga cct act gca Tyr Val Tyr Asn Val Gly Asp Arg Glu Gln Leu Asp Arg Pro Thr Ala 100 105 110	573
aaa ctc atc aag gat ttt tct ttg ata gaa gat tct ttt atg agc acc Lys Leu Ile Lys Asp Phe Ser Leu Ile Glu Asp Ser Phe Met Ser Thr 115 120 125 130	621
att gga agt cga tat gga ata aca gct gga tct cgg gag aat acc ctg Ile Gly Ser Arg Tyr Gly Ile Thr Ala Gly Ser Arg Glu Asn Thr Leu 135 140 145	669
tat aat gct ctt tgg gtt gct cag gcg ctg ttg cgc aag gga tct gtg Tyr Asn Ala Leu Trp Val Ala Gln Ala Leu Leu Arg Lys Gly Ser Val 150 155 160	717
aag acc gtg agt aaa cga att ctt ata ttc acc aat gaa gat gat cct Lys Thr Val Ser Lys Arg Ile Leu Ile Phe Thr Asn Glu Asp Asp Pro 165 170 175	765
ttt ggt act att aca gga gca gtg aaa act gat atg att agg aca aca Phe Gly Thr Ile Thr Gly Ala Val Lys Thr Asp Met Ile Arg Thr Thr 180 185 190	813
ctt caa cgt gca aag gat gca caa gat ctt ggc ctg tct att gaa ctt Leu Gln Arg Ala Lys Asp Ala Gln Asp Leu Gly Leu Ser Ile Glu Leu 195 200 205 210	861
ctt cca ctg agc ccg cct gat gac cag ttc aac atg tcc ctg ttt tat Leu Pro Leu Ser Pro Pro Asp Asp Gln Phe Asn Met Ser Leu Phe Tyr 215 220 225	909
gca gat ttg att ggt ctg gat gga gat gag atg acc gaa tat ttg cca Ala Asp Leu Ile Gly Leu Asp Gly Asp Glu Met Thr Glu Tyr Leu Pro 230 235 240	957
tct gct ggt gat aag cta gag gat atg act aat caa ctg aga aaa cgg Ser Ala Gly Asp Lys Leu Glu Asp Met Thr Asn Gln Leu Arg Lys Arg 245 250 255	1005
ata atg aag aag cgc aga atc aaa act ctt tca ttt gcg att acc aat Ile Met Lys Lys Arg Arg Ile Lys Thr Leu Ser Phe Ala Ile Thr Asn 260 265 270	1053
gat gtg tgc ata gaa gtc aat aca tat gcg ctg gtc cgt cct act act Asp Val Cys Ile Glu Val Asn Thr Tyr Ala Leu Val Arg Pro Thr Thr 275 280 285 290	1101
aca ggg aca atc aca tgg ctt gat tca cta agt aac ctc cca tta aag Thr Gly Thr Ile Thr Trp Leu Asp Ser Leu Ser Asn Leu Pro Leu Lys 295 300 305	1149
gtt gag agg tct ttc ata tgc aat gat act ggg gct cta ctt cag gat Val Glu Arg Ser Phe Ile Cys Asn Asp Thr Gly Ala Leu Leu Gln Asp 310 315 320	1197

-3-

gca cag aca cgt ttc cag atg tac aat gac aca att gtc aaa ttt tct Ala Gln Thr Arg Phe Gln Met Tyr Asn Asp Thr Ile Val Lys Phe Ser 325 330 335	1245
gta cgt gaa ctc tct gag gtt aaa agg gtt gca agc cat cat ctt cgc Val Arg Glu Leu Ser Glu Val Lys Arg Val Ala Ser His His Leu Arg 340 345 350	1293
ctt ata ggt ttc aag cca ttg gat tgc ttg aaa gat tac cat aac tta Leu Ile Gly Phe Lys Pro Leu Asp Cys Leu Lys Asp Tyr His Asn Leu 355 360 365 370	1341
aga cca tcg aca ttt att tat ccg agt gat gag cgt ata ttt gga agc Arg Pro Ser Thr Phe Ile Tyr Pro Ser Asp Glu Arg Ile Phe Gly Ser 375 380 385	1389
acc tgt gtt ttc gtt gct tta cat agc tca atg tta cgt ctt gga agg Thr Cys Val Phe Val Ala Leu His Ser Ser Met Leu Arg Leu Gly Arg 390 395 400	1437
ttt gca ctt gca ttt tat ggg aat cca act cga cca caa ctc ata gcc Phe Ala Leu Ala Phe Tyr Gly Asn Pro Thr Arg Pro Gln Leu Ile Ala 405 410 415	1485
ctt gtt gct caa gaa gag gtt act tcg tct ggt cgt cag ttt gaa ccg Leu Val Ala Gln Glu Glu Val Thr Ser Ser Gly Arg Gln Phe Glu Pro 420 425 430	1533
cct ggc atg cac atg atc tat ctt cca tac tcc gat gat att aga tat Pro Gly Met His Met Ile Tyr Leu Pro Tyr Ser Asp Asp Ile Arg Tyr 435 440 445 450	1581
cct gaa gaa gta agg act aac gtt gta ttt caa ctt tac gat ctc tgt Pro Glu Glu Val Arg Thr Asn Val Val Phe Gln Leu Tyr Asp Leu Cys 455 460 465	1629
aag cta att ttc ttg ctt ggt ttg cag gtt cat gtg act tct gat gat Lys Leu Ile Phe Leu Leu Gly Leu Gln Val His Val Thr Ser Asp Asp 470 475 480	1677
gca ccg cgt gca aca gat gaa caa atc aag aaa gct tcg aac ata ttc Ala Pro Arg Ala Thr Asp Glu Gln Ile Lys Lys Ala Ser Asn Ile Phe 485 490 495	1725
aaa cgt att gat ctg ata aat ttc tct gca tgc caa ttt gct aac cca Lys Arg Ile Asp Leu Ile Asn Phe Ser Ala Cys Gln Phe Ala Asn Pro 500 505 510	1773
gct ttg caa aga cac tat ggg atc ttg gag gcc tta gct tta ggc gaa Ala Leu Gln Arg His Tyr Gly Ile Leu Glu Ala Leu Ala Leu Gly Glu 515 520 525 530	1821
tat gag atg cct gat ata aag gat gag acc ctg cct gac gaa gaa ggc Tyr Glu Met Pro Asp Ile Lys Asp Glu Thr Leu Pro Asp Glu Glu Gly 535 540 545	1869
ttg tct aag cca ggg gta gcc aat gct att gag gaa ttc aag act tca	1917

-4-

Leu Ser Lys Pro Gly Val Ala Asn Ala Ile Glu Glu Phe Lys Thr Ser
 550 555 560
 gtc tat ggt gaa aat tat gac caa gag gag gca gaa gcg gca gca ggg 1965
 Val Tyr Gly Glu Asn Tyr Asp Gln Glu Glu Ala Glu Ala Ala Ala Gly
 565 570 575
 aaa gct tcc cgt ggt aat gct tca aaa aag cgg aag gag gtc act gat 2013
 Lys Ala Ser Arg Gly Asn Ala Ser Lys Lys Arg Lys Glu Val Thr Asp
 580 585 590
 gca gct gcg cag ata agt gct gct tat gat tgg gca gaa ctt gca gac 2061
 Ala Ala Ala Gln Ile Ser Ala Ala Tyr Asp Trp Ala Glu Leu Ala Asp
 595 600 605 610
 aat gga aaa ctg aag gaa atg acc acg gtg gaa ttg aga tcc tac ctg 2109
 Asn Gly Lys Leu Lys Glu Met Thr Thr Val Glu Leu Arg Ser Tyr Leu
 615 620 625
 acc gcg cat gat ctc ccg gtt tct ggt aag aaa gat gta ctt atc agc 2157
 Thr Ala His Asp Leu Pro Val Ser Gly Lys Lys Asp Val Leu Ile Ser
 630 635 640
 agg atc ttg act cac ctg ggt aag tgaagccggg tatccgaact gttagtttcc 2211
 Arg Ile Leu Thr His Leu Gly Lys
 645 650
 tgagaccgga ctgacatctg tagtgctgtc tcgcggatcc ttgtggaccg taatectccc 2271
 ccgtgcaggc tacgtgccga ttttttgtat agcgacagga cagcaccagt tgccaatgta 2331
 acctgtgtta agtttctgtc tgtggtaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2391
 aaaaaaaaaa 2399

<210> 2
 <211> 650
 <212> PRT
 <213> Zea mays

<400> 2
 Met Asp Leu Asp Pro Glu Gly Ile Phe Arg Asp Asp Ser Asp Glu Asp
 1 5 10 15
 Glu Asp Ser Val Gln Glu Arg Glu Ala Asn Lys Glu Met Val Val Tyr
 20 25 30
 Leu Val Asp Ala Ser Pro Lys Met Phe Thr Pro Ala Thr Thr Gln Asp
 35 40 45
 Asn Glu Lys Gln Glu Thr His Phe His Thr Ile Val Asn Cys Ile Thr
 50 55 60
 Glu Ser Leu Lys Thr Gln Ile Ile Gly Arg Ser Tyr Asp Glu Val Ala
 65 70 75 80
 Ile Cys Phe Phe Asn Thr Lys Glu Lys Lys Asn Leu Gln Asp Ser Ala
 85 90 95
 Gly Val Tyr Val Tyr Asn Val Gly Asp Arg Glu Gln Leu Asp Arg Pro
 100 105 110
 Thr Ala Lys Leu Ile Lys Asp Phe Ser Leu Ile Glu Asp Ser Phe Met
 115 120 125
 Ser Thr Ile Gly Ser Arg Tyr Gly Ile Thr Ala Gly Ser Arg Glu Asn
 130 135 140
 Thr Leu Tyr Asn Ala Leu Trp Val Ala Gln Ala Leu Leu Arg Lys Gly

-5-

145					150					155				160
Ser	Val	Lys	Thr	Val	Ser	Lys	Arg	Ile	Leu	Ile	Phe	Thr	Asn	Glu
				165					170					175
Asp	Pro	Phe	Gly	Thr	Ile	Thr	Gly	Ala	Val	Lys	Thr	Asp	Met	Ile
			180				185						190	
Thr	Thr	Leu	Gln	Arg	Ala	Lys	Asp	Ala	Gln	Asp	Leu	Gly	Leu	Ser
		195					200					205		Ile
Glu	Leu	Leu	Pro	Leu	Ser	Pro	Pro	Asp	Asp	Gln	Phe	Asn	Met	Ser
	210					215				220				Leu
Phe	Tyr	Ala	Asp	Leu	Ile	Gly	Leu	Asp	Gly	Asp	Glu	Met	Thr	Glu
225					230				235					240
Leu	Pro	Ser	Ala	Gly	Asp	Lys	Leu	Glu	Asp	Met	Thr	Asn	Gln	Leu
			245						250					255
Lys	Arg	Ile	Met	Lys	Lys	Arg	Arg	Ile	Lys	Thr	Leu	Ser	Phe	Ala
		260					265						270	Ile
Thr	Asn	Asp	Val	Cys	Ile	Glu	Val	Asn	Thr	Tyr	Ala	Leu	Val	Arg
	275					280					285			Pro
Thr	Thr	Thr	Gly	Thr	Ile	Thr	Trp	Leu	Asp	Ser	Leu	Ser	Asn	Leu
	290				295						300			Pro
Leu	Lys	Val	Glu	Arg	Ser	Phe	Ile	Cys	Asn	Asp	Thr	Gly	Ala	Leu
305					310				315					320
Gln	Asp	Ala	Gln	Thr	Arg	Phe	Gln	Met	Tyr	Asn	Asp	Thr	Ile	Val
		325						330						335
Phe	Ser	Val	Arg	Glu	Leu	Ser	Glu	Val	Lys	Arg	Val	Ala	Ser	His
		340					345					350		His
Leu	Arg	Leu	Ile	Gly	Phe	Lys	Pro	Leu	Asp	Cys	Leu	Lys	Asp	Tyr
	355					360					365			His
Asn	Leu	Arg	Pro	Ser	Thr	Phe	Ile	Tyr	Pro	Ser	Asp	Glu	Arg	Ile
	370				375						380			Phe
Gly	Ser	Thr	Cys	Val	Phe	Val	Ala	Leu	His	Ser	Ser	Met	Leu	Arg
385				390					395					400
Gly	Arg	Phe	Ala	Leu	Ala	Phe	Tyr	Gly	Asn	Pro	Thr	Arg	Pro	Gln
		405						410						415
Ile	Ala	Leu	Val	Ala	Gln	Glu	Glu	Val	Thr	Ser	Ser	Gly	Arg	Gln
		420					425					430		Phe
Glu	Pro	Pro	Gly	Met	His	Met	Ile	Tyr	Leu	Pro	Tyr	Ser	Asp	Asp
	435					440					445			Ile
Arg	Tyr	Pro	Glu	Glu	Val	Arg	Thr	Asn	Val	Val	Phe	Gln	Leu	Tyr
	450				455						460			Asp
Leu	Cys	Lys	Leu	Ile	Phe	Leu	Leu	Gly	Leu	Gln	Val	His	Val	Thr
465				470					475					480
Asp	Asp	Ala	Pro	Arg	Ala	Thr	Asp	Glu	Gln	Ile	Lys	Lys	Ala	Ser
		485						490					495	Asn
Ile	Phe	Lys	Arg	Ile	Asp	Leu	Ile	Asn	Phe	Ser	Ala	Cys	Gln	Phe
	500						505						510	Ala
Asn	Pro	Ala	Leu	Gln	Arg	His	Tyr	Gly	Ile	Leu	Glu	Ala	Leu	Ala
	515					520					525			Leu
Gly	Glu	Tyr	Glu	Met	Pro	Asp	Ile	Lys	Asp	Glu	Thr	Leu	Pro	Asp
	530					535				540				Glu
Glu	Gly	Leu	Ser	Lys	Pro	Gly	Val	Ala	Asn	Ala	Ile	Glu	Glu	Phe
545				550					555					Lys
Thr	Ser	Val	Tyr	Gly	Glu	Asn	Tyr	Asp	Gln	Glu	Glu	Ala	Glu	Ala
		565						570					575	Ala
Ala	Gly	Lys	Ala	Ser	Arg	Gly	Asn	Ala	Ser	Lys	Lys	Arg	Lys	Glu
	580						585					590		Val
Thr	Asp	Ala	Ala	Ala	Gln	Ile	Ser	Ala	Ala	Tyr	Asp	Trp	Ala	Glu
	595					600						605		Leu

-6-

Ala Asp Asn Gly Lys Leu Lys Glu Met Thr Thr Val Glu Leu Arg Ser
610 615 620
Tyr Leu Thr Ala His Asp Leu Pro Val Ser Gly Lys Lys Asp Val Leu
625 630 635 640
Ile Ser Arg Ile Leu Thr His Leu Gly Lys
645 650

<210> 3

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide based upon an adaptor
used for cDNA library construction and poly(dT) to
remove clones which have a poly(A) tail but no
cDNA insert.

<400> 3

tcgacccacg cgtccgaaaa aaaaaaaaaa aaaaaa

36